

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/00	A2	(11) International Publication Number: WO 99/47557 (43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/US99/06051 (22) International Filing Date: 18 March 1999 (18.03.99) (30) Priority Data: 60/078,634 18 March 1998 (18.03.98) US (71) Applicant: ONYX PHARMACEUTICALS, INC. [US/US]; 3031 Research Drive, Richmond, CA 94806 (US). (72) Inventors: BOLLAG, Gideon; 172 Catalina Drive, Hercules, CA 94547 (US). HART, Matthew, J.; 1210 Masonic Avenue, Berkeley, CA 94706 (US). ROSCOE, William; 3099 California Street, San Francisco, CA 94115 (US). POLAKIS, Paul; 509 Barone Lane, Mill Valley, CA 94941 (US). STERNWEIS, Paul; 2103 Flat Creet Drive, Richardson, TX 75080 (US). KOZASA, Tohru; 7415 Centenary, Dallas, TX 75225 (US). JIANG, Xuejun; 2400 Waterview Parkway #932, Richardson, TX 75080 (US). (74) Agent: GIOTTA, Gregory; Onyx Pharmaceuticals, Inc., 3031 Research Drive, Richmond, CA 94806 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: IDENTIFICATION OF FACTORS WHICH MEDIATE THE INTERACTION OF HETEROTRIMERIC G PROTEINS AND MONOMERIC G PROTEINS (57) Abstract Monomeric GTPase guanine nucleotide exchange factor (GEF) have been identified which also contain an RGS region analogous to those of GTPase activating proteins (GAP). One of these GEF proteins, a Rho GEF has been demonstrated to contain an RGS sequence that has GAP activity toward a α subunit of a heterotrimeric G prote in.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

IDENTIFICATION OF FACTORS WHICH MEDIATE THE INTERACTION OF HETEROTRIMERIC G PROTEINS AND MONOMERIC G PROTEINS

BACKGROUND OF THE INVENTION

5 Signal transduction pathways linking extracellular factors to the activation of the Rho GTPase have been implicated in cell growth control and cytoskeletal rearrangements. Specifically, heterotrimeric G proteins have been shown to mediate these pathways, although the mechanism of mediation has been unclear. The identification of factors which interact with both heterotrimeric G proteins and Rho GTPase would provide an important
10 tool for investigating and controlling various cell processes, including cell proliferative diseases.

SUMMARY OF THE INVENTION

 The invention relates to a polypeptide, and corresponding nucleic acid, comprising an amino acid sequence of a novel RGS domain, obtainable, e.g., from a guanine nucleotide
15 exchange factor (GEF) protein, where the polypeptide preferably does not include a dbl homology (DH) domain or a pleckstrin homology (PH) domain. In a preferred embodiment, the polypeptide has GTPase activating activity and binding affinity for an α G protein subunit such as $G\alpha$.

 The polypeptides and nucleic acids can be used as tools for research, therapeutics,
20 and diagnostics as discussed below.

 The invention also relates to a method of identifying or assaying for a molecule, or mixture of molecules, that regulate the binding of an RGS domain of a GEF protein to a substrate, e.g., a G protein subunit such as $G\alpha$. In one embodiment, the method involves incubating, under effective conditions, a polypeptide having an RGS domain of a GEF
25 polypeptide, and optionally having GEF activity, with a $G\alpha$ subunit, or a fragment thereof, in the presence and/or absence of a test molecule; and determining whether the presence of the test molecule regulates the binding between the polypeptide and the subunit, or fragment thereof. As discussed later, various RGS-GEF polypeptides binding substrates can be utilized.

30 In addition, the invention relates to a method of identifying or assaying for a molecule, or mixture of molecules, that regulates a stimulatory effect of a polypeptide

comprising an RGS domain of a GEF protein on a polypeptide having a GTPase activity. In a preferred embodiment, the method comprises incubating a $G\alpha$ subunit and a GEF protein, under effective conditions, in the presence and absence of a test molecule and determining whether the presence of the test molecule regulates the stimulatory effect of the GEF protein on $G\alpha$ subunit GTPase activity.

The invention also relates to a method of identifying or assaying for a molecule that specifically regulates a stimulatory effect of a first polypeptide, such as an activated $G\alpha$ subunit, or polypeptide having GTPase activity, on a nucleotide exchange factor activity of a second polypeptide. The second polypeptide preferably comprises a RGS-GEF domain obtainable from a GEF, and more preferably is a guanine nucleotide exchange factor (GEF) for a monomeric G protein. In one embodiment of the method, a first assay is conducted by incubating an activated $G\alpha$ subunit with a GEF protein and a monomeric G protein in the presence and absence of a test molecule; a second assay is conducted: by incubating a GEF protein and a monomeric G protein in the presence and absence of the test molecule, and a determination is made as to whether the molecule has a different effect when the first assay is compared to the second assay.

The invention further relates to a method of identifying or assaying for a molecule, or mixture of molecules, that mimics the stimulatory effect of an activated $G\alpha$ subunit on GEF mediated nucleotide exchange of a monomeric G protein. In one example, such a method comprises identifying a test compound that exhibits a binding affinity for an RGS domain of GEF proteins, incubating a GEF protein and monomeric G protein in the presence or absence of the test compound, determining whether the test compound exhibits a stimulatory effect on GEF mediated nucleotide exchange of a monomeric G protein.

The invention further relates to a method of identifying or assaying for a molecule, or mixture of molecules, that mimics the stimulatory effect of an RGS domain of GEF polypeptide on $G\alpha$ subunit GTPase activity. In one example, such a method comprises identifying a test compound that exhibits a binding affinity for a $G\alpha$ subunit and incubating a GTP loaded $G\alpha$ subunit in the presence or absence of the test compound to determine whether the test compound exhibits a stimulatory effect on GEF mediated nucleotide exchange of a monomeric G protein.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1, Panel A depicts the alignment of the sequences from RGS proteins and the N-terminal region of p115 Rho GEF as performed by Clustal W with a secondary structure mask of RGS4 to assign penalties for gaps. The RGS homologous sequences of Lsc, KIAA380, and DrhoGEF2 were further added to this alignment by Clustal W and manual adjustments. The (a) symbols above RGS4 indicate the α helices of the RGS domain of RGS4. Dark shaded boxes indicate conserved residues of the hydrophobic core of the RGS structure. Lightly shaded boxes show other conserved residues. Asterisks mark the residues of RGS4 which contact $G\alpha_{i1}$. Primary sequences used in the alignment are the following:

10 rat RGS4 (SwissProt accession number P49799), mouse RGS2 (O08849), human GAIIP (P49795), rat RGS12 (O08774), rat RGS14 (O08773), human p115 (1654344), mouse Lsc (1389756), human KIAA380 (2224701) and Drosophila DrhoGEF2 (2760368).

Figure 1, Panel B depicts constructs of p115 Rho GEF that were employed in the studies described herein. Numbers indicate the residues of p115 in each construct. The RGS, dbl(DH), and pleckstrin (PH) homology regions are indicated. GST=glutathione-S-transferase.

15

Figure 2, Panel A is a graph showing the hydrolysis of GTP bound to $G\alpha_{i3}$ and $G\alpha_{i2}$ at 15°C either with (●○) or without (■□) 10 nM p115 Rho GEF.

Figure 2, Panel B is a graph showing the hydrolysis at 4°C of GTP bound to $G\alpha_{i3}$ (●) and $G\alpha_{i2}$ (○) and in the presence of various concentrations of p115 Rho GEF. The initial rates of reaction were plotted as a function of the concentration of p115 Rho GEF.

20

Figure 3 is a graph showing the hydrolysis at 15°C of GTP bound to $G\alpha_{i3}$ and $G\alpha_{i2}$ with either full-length p115 Rho GEF (●), $\Delta Np115$ (■), or RGS-p115 (▲), or without any p115 construct (▼).

25

Figure 4 is a graph showing the hydrolysis of GTP bound to $G\alpha_{i1}$, $G\alpha_z$, $G\alpha_q$, and $G\alpha_s$ with 100 nM p115 Rho GEF (Δ), 100 nM RGS4 (\square), or buffer control (○). Assays were performed at 4°C for $G\alpha_{i1}$ and $G\alpha_s$, at 15°C for $G\alpha_z$, and at 20°C for $G\alpha_q$.

Figure 5 is a graph showing the selective inhibition of p115 GAP activity by the AlF_4^- -activated forms of $G\alpha$ subunits. Panel A: P115 (400 nM) was incubated on ice for 15 minutes with various $G\alpha$ subunits (400 nM) in the presence of 30 μM $AlCl_3$, 10 mM NaF,

30

and 10 mM MgSO₄. The mixture was diluted 20-fold, mixed with 0.3 nM Gα₁₂(GTP) and the hydrolysis of bound GTP was measured after incubation at 15°C for 2 minutes. Panel B: P115 (400 nM) was incubated with various concentrations of Gα₁₂(GDP-AIF₄⁻) (●) or Gα₁₃(GDP-AIF₄⁻) (■) as described for Panel A. The mixture was diluted 20-fold, mixed with 1nM Gα₁₃(GTP) at 4°C and the hydrolysis of bound GTP was assessed over time. The initial rate of GTPase of Gα₁₃ was plotted against the final concentration of α subunit GDP-AIF₄⁻. The filled triangle indicates the rate of GTPase of Gα₁₃ without p115.

Figure 6, Panel A is an image of an immunoblot showing the detection of myc-tagged p115 Rho GEF expression in COS cells using an anti-myc antibody.

Figure 6, Panel B is an image of an immunoblot showing the detection of a coimmunoprecipitate of p115 Rho GEF and Gα₁₃ using an anti-myc antibody.

Figure 6, Panel C is an image of an immunoblot showing the detection of the coimmunoprecipitate of p115 Rho GEF and Gα₁₃ using an anti-Gα₁₃ antibody.

Figure 6, Panel D is an image of an immunoblot showing the detection of p115 Rho GEF and Gα₁₃ binding when purified Gα₁₃ is added to immunoprecipitated p115 Rho GEF when using an anti Gα₁₃ antibody.

Figure 7, Panel A is a graph showing the dissociation of bound GDP from 100 nM RhoA after 10 minutes in the presence or absence of 100 nM Gα₁₃ or Gα₁₂ and in the presence of various concentrations of p115 Rho GEF as indicated.

Figure 7, Panel B is a graph showing the dissociation of GDP from 100 nM RhoA after 10 minutes in the presence of 25 nM p115 Rho GEF and the indicated concentrations of Gα₁₃ or Gα₁₂. Unstimulated dissociation of GDP from RhoA is indicated by the lower dashed line.

Figure 7, Panel C is a graph showing the dissociation of GDP from 100 nM RhoA after 10 minutes of incubation with p115 Rho GEF and Gα₁₃ that had been treated with AMF, GTPγS or GDPβS as indicated.

Figure 7, Panel D is a graph showing the dissociation of of GDP from 100 nM RhoA after 10 minutes of incubation with p115 Rho GEF (25 nM) and various Gα subunits (100 nM) as indicated.

Figure 8, Panel A is a graph showing the association of 1 nM [32 P]GTP to 100 nM RhoA in the presence of the indicated concentrations of truncated for full-length p115 Rho GEF as measured by filtration after 30 minutes at 30°C.

Figure 8, Panel B is a graph showing the dissociation of [3 H]-GDP from 100 nM RhoA after incubation for 10 minutes in the presence or absence of 25 nM p115 Rho GEF, 20 nM $G\alpha_{13}$, and 300 nM GST-RGSp115 as indicated.

Figure 8, Panel C is a graph showing the dissociation of [3 H]-GDP from 100 nM RhoA after incubation for 10 minutes in the presence 25 nM p115 Rho GEF and in the presence or absence of 25 nM $G\alpha_{13}$ and the indicated concentrations of $G\alpha_{12}$.

Figure 9, Panel A is an image of an immunoblot showing the detection of myc-tagged KIAA380 (designated FL147) expression in COS cells using an anti-myc antibody.

Figure 9, Panel B is an image of an immunoblot showing the detection of a coimmunoprecipitate of KIAA380 (designated FL147) and $G\alpha_{12}$ using an anti- $G\alpha_{12}$ antibody.

Figure 10 is the a listing of the amino acid sequence for p115 Rho GEF.. The RGS domain is shown by amino acids 45-170.

Figure 11 is a listing of the nucleic acid sequence for p115 Rho GEF. The RGS domain is encoded by nucleotides 187-564.

Figure 12 is a listing of the amino acid sequence for KIAA380. The RGS domain is shown by amino acids 310-432.

Figure 13 is a listing of the nucleic acid sequence for KIAA380. The RGS domain is encoded by nucleotides 1673-2041.

Figure 14 is a listing of the amino acid sequence for Lsc. The RGS domain is shown by amino acids 43-168.

Figure 15 is a listing of the nucleic acid sequence for Lsc. The RGS domain is encoded by nucleotides 218-595.

Figure 16 is a listing of the amino acid sequence for DRhoGEF2. The RGS domain is shown by amino acids 924-1053

Figure 17 is a listing of the nucleic acid sequence for DRhoGEF2. The RGS domain is encoded by nucleotides 3185-3574.

Figure 18 is a homology alignment of the RGS region of several proteins, including GEF proteins with RGS domains (e.g. p115 Rho GEF, Lsc, KIAA380, DrhoGEF). The alignment was performed using the Clustal method with a PAM250 residue weight table.

DETAILED DESCRIPTION OF THE INVENTION

5 G proteins transduce signals from a large number of cell surface heptahelical receptors to various intracellular effectors. Each heterotrimeric G protein is composed of a guanine nucleotide-binding α subunit and a high-affinity dimer of β and γ subunits. $G\alpha$ subunits are commonly classified into four subfamilies (G_s , G_i , G_q , and G_{12}) based on their amino acid sequence homology and function (A.G. Gilman, *Annu. Rev. Biochem.*, **56**, 615
10 (1987); Y. Kaziro et al., *Annu. Rev. Biochem.*, **60**, 349 (1991); Hepler and Gilman, *Trends Biochem. Sci.*, **17**, 383, (1992)). The G_{12} subfamily, consists of two identified members to date, G_{12} and G_{13} .

In accordance with the present invention, the identification of proteins having activity as both a GTPase activating protein (GAP) for the α subunit of a heterotrimeric G
15 protein and activity as a guanine nucleotide exchange factor (GEF) activity for monomeric G proteins have been described. Also in accordance with the invention, the first identification of a protein having GAP activity for the G_{12} subfamily of G proteins has been described. Also in accordance with the invention, the ability of an α subunit of a heterotrimeric G protein to stimulate GEF mediated guanine nucleotide exchange activity of a monomeric G
20 protein has been described. GAP and GEF activity, and methods of screening thereof, are described in Berman et al., 1996, *Cell* **86**:445 and Hart et al., 1996, *J. Biol. Chem.*, **271**:25452.

According to the present invention, the GAP activity of GEF proteins has been correlated with a novel RGS domain obtainable from a GEF protein. The present invention
25 relates to all aspects of such an RGS domain, including all aspects of a Rho GEF such as p115 Rho-GEF. (U.S. Patent Application No. 08/943,768, herein incorporated by reference).

A GEF protein modulates cell signaling pathways, both in *in vitro* and *in vivo*, by modulating the guanine nucleotide exchange activity of a GTPase. According to the present invention, a GEF protein which also modulates the GTPase activity of a heterotrimeric $G\alpha$
30 subunit is described. By way of illustration, p115 Rho-GEF, which modulates the guanine

nucleotide exchange activity of a Rho GTPase, as well as the GTPase activity of the $G\alpha_{12}$ family of heterotrimeric G protein subunits is described.

The present invention particularly relates to polypeptides comprising a RGS domain of a GEF polypeptide, or fragments thereof, and corresponding nucleic acids.

5 The invention also relates to methods of using such polypeptides, nucleic acids, or derivatives thereof, e.g., in therapeutics, diagnostics, and as research tools. Other aspects of the present invention relate to antibodies and other ligands which recognize the RGS domain of GEF polypeptides or nucleic acids, methods for identifying or assaying modulators of the GEF activities and/or the GAP activities of a protein containing a RGS domain, and
10 methods of treating pathological conditions associated with or related to the RGS domain, e.g., a GEF mediated interaction of a $G\alpha$ subunit and a Rho GTPase.

As used herein, an "RGS-GEF polypeptide" means, e.g., a polypeptide containing an RGS domain derived from a GEF protein, such as p115 Rho-GEF, Lsc, KIAA0380, or DRhoGEF2, and, which has one or more of the following activities: a specific binding
15 affinity for a polypeptide substrate, e.g., a G protein subunit, preferably an α subunit, such as G_{12} or G_{13} ; a GTPase activating activity (GAP), such as a GAP activity for a G protein α subunit; or, an immunogenic activity. An RGS-GEF polypeptide preferably does not contain a (dbl homology) DH or a (pleckstrin homology) PH domain. DH and PH domains are disclosed in Cerione and Zheng, 1996, *Curr. Opin. In Cell Biol.*, 8:216. For example, the
20 amino acid sequence of p115 Rho GEF (Fig. 10) contains a novel RGS domain at amino acids 45-170, the DH domain at amino acids 420-637, and the PH domain at amino acids 646-672. By "derived," it is meant that the amino acid sequence is obtainable from a naturally-occurring GEF (such as p115, Lsc, KIAA380, and DrrhoGEF2) or a non-naturally-occurring "mutated" sequence which is based upon a naturally-occurring GEF sequence
25 (i.e., different amino acid residues have been substituted for the amino acid residues which occur in the naturally-occurring sequence at a particular position). The polypeptide can be "isolated," i.e., the material is in a form in which it is not found in its original environment, e.g., more concentrated, more purified, or separated from other components, etc. A preferred RGS polypeptide possesses both a GAP and GEF activity, e.g., a mutated p115 Rho-GEF.
30 See below.

An RGS-GEF nucleic acid codes for an RGS-GEF polypeptide. The nucleic acid refers to both sense and anti-sense nucleic acids.

By the term "specific binding affinity," it is meant, e.g., that the RGS-GEF polypeptide has a binding preference for the activated state or transition state of a G protein subunit as compared to a GDP-bound state or the nucleotide depleted state. By "GEF activity," it is meant, e.g., that the polypeptide stimulates or catalyzes the dissociation of GDP from a monomeric G-protein, such as Rho, and subsequent binding of GTP. Monomeric G-proteins include but are not limited to G-proteins in the Ras, Rho/Rac, Sar, Rab, Arf, and Ran families. Of particular interest are the RGS domains of the following GEF proteins: human p115 (1654344) (Fig. 10, RGS domain at amino acids 45-170), mouse Lsc (1389756) (Fig. 14, RGS domain at amino acids 43-168), KIAA380 (2224701) (Fig. 12, RGS domain at amino acids 310-432) and Drosophila DrhoGEF2 (2760368) (Fig. 16, RGS domain at amino acids 924-1053).

Another aspect of the invention relates to novel consensus sequences for RGS domain(s) of a GEF protein, herein referred to as a "sub-RGS consensus sequence." An "RGS domain," as used herein, refers to the amino acid sequence of protein which is able to bind to or physically interact with a G protein and, optionally, stimulates GTPase activity of that protein. A "sub-RGS consensus sequence," as used herein, refers to a consensus sequence which can be used to identify a specific subset of proteins which contain an RGS domain. For example, a homology alignment of the RGS domain from several proteins as shown and described in Fig. 18 and the corresponding legend, shows that several sub-RGS consensus sequences may be defined by the gap of 13 to 14 amino acids that is apparent in the RGS domains of GEF proteins. One of these consensus sequences, herein designated as "RGS-GEF consensus 1," is herein defined to be a consensus sequence of AA₁-AA₂-AA₃-AA₄-AA₅-AA₆-AA₇-AA₈-(gap of 13 amino acids)-AA₂₂-AA₂₃-AA₂₄-AA₂₅-AA₂₆, wherein:

- AA₁ is L;
- AA₂ is E or V;
- AA₃ is K or P;
- AA₄ is T, N, or R;
- AA₅ is A;
- AA₆ is V or P

AA₇ is L;

AA₈ is either S or a gap of one amino acid, contiguous with the gap of 13 amino acids;

AA₂₂ is either R or W;

AA₂₃ is either V or Y;

5 AA₂₄ is either P, K, or R

AA₂₅ is either V, I, or Q;

AA₂₆ is either P or D.

A second consensus sequence, herein designated as "RGS-GEF consensus 2," is herein defined to be a consensus sequence of AA₁-AA₂-AA₃-AA₄-(gap of 13 amino acids)-

10 AA₁₈-AA₁₉, wherein:

AA₁ is A;

AA₂ is V or P;

AA₃ is L;

AA₄ is either S or a gap of one amino acid, contiguous with the gap of 13 amino acids;

15 AA₁₈ is either R or W;

AA₁₉ is either V or Y.

Other proteins, including other GEF proteins can be aligned with the RGS domain of RGS proteins as shown in Figure 18, and using methods described herein, to determine if they contain a sub-RGS consensus sequence, such as RGS-GEF consensus 1 or RGS-GEF

20 consensus 2, as defined above.

In examining Figure 18 it is also apparent that a nucleotide sequence unique to RGS proteins that are not GEF proteins is shown by the nucleotide sequences which encode the amino acids that correspond to the 13-14 amino acid gap in RGS-GEF proteins. These nucleotide sequences could be used as probes to identify particular types of RGS proteins.

25 RGS-GEF polypeptides are preferably biologically-active. By biologically-active, it is meant that a polypeptide fragment possesses an activity in a living system or with component(s) of a living system. Biological-activities include, but are not limited to a specific binding affinity for a G protein α subunit, as defined above, and GAP activity toward a G protein α subunit. As described in the examples, such polypeptides can be

30 prepared routinely, e.g., by recombinant means or by proteolytic cleavage of isolated polypeptides, and then assayed for a desired activity.

A polypeptide of the present invention includes polypeptides which have less than 100% identity to the amino acid sequences of p115 Rho-GEF (Fig. 10), Lsc (Fig. 14), KIAA0380 (Fig. 12), or DRhoGEF2 (Fig. 16). For the purposes of the following discussion: Sequence identity means that the same nucleotide or amino acid which is found in the sequences set forth in Fig. 10-17 is found at the corresponding position of the compared sequence(s). A polypeptide having less than 100% sequence identity to the amino acid sequences set forth in Figures 10, 12, 14, and 16 can be substituted in various ways, e.g., by a conservative amino acid. The sum of the identical and conservatively substituted residues divided by the total number of residues in the sequence is equal to the percent sequence similarity. For purposes of calculating sequence identity and similarity, the compared sequences can be aligned and calculated according to any desired method, algorithm, computer program, etc., including, e.g., FASTA, BLASTA. A polypeptide having less than 100% amino acid sequence identity to the amino acid sequences of the GEF proteins shown in Figures 10, 12, 14, and 16 may comprise, for example, about 60, 65 percent sequence similarity and more preferably about 67, 70, 78, 80, 90, 92, 96, 99, etc. percent sequence amino acid sequence similarity.

In particular, the present invention relates to polypeptides, and corresponding nucleic acids, of p115, Lsc, KIAA380, and DrhoGEF2 which are mutated in the RGS domain of a GEF protein and which possess one or more of the RGS-GEF polypeptide activities mentioned above. By the term "mutated," it is meant herein that such sequences are not naturally-occurring. For example a mutated polypeptide as mentioned can have one or more naturally-occurring positions replaced by a conservative amino acid, e.g., (based on the size of the side chain and degree of polarization) small nonpolar: cysteine, proline, alanine, threonine; small polar: serine, glycine, aspartate, asparagine; large polar: glutamate, glutamine, lysine, arginine; intermediate polarity: tyrosine, histidine, tryptophan; large nonpolar: phenylalanine, methionine, leucine, isoleucine, valine. Such conservative substitutions also include those described by Dayhoff in the Atlas of Protein Sequence and Structure 5 (1978), and by Argos in EMBO J., 8, 779-785 (1989). A polypeptide having an amino acid sequence as set forth in Figures 10, 12, 14, and 16 can be substituted at 1, 5, 10, 15, or 20 positions by conservative amino acids. The mutations can be introduced into the conserved consensus region or the other residues of the RGS domain of a GEF protein.

A mutation to an RGS-GEF polypeptide can be selected to have one or more of the activities mentioned above, e.g., a specific binding affinity for a G protein α subunit, a GAP activity toward a G protein α subunit, etc.. Assays for such activities can be conducted as described below or as disclosed in Cerione and Zheng, 1996, *Curr. Opin. In Cell Biol.*, 8:216.

An RGS-GEF polypeptide can be modified by introducing amino acid substitutions into the hydrophobic core of the RGS domain (See Fig. 1, Panel A). For example, a conservative amino acid substitution would not be expected to affect activity, whereas as non-conservative amino acid substitution, e.g., changing a hydrophobic residue to a hydrophilic residue, would be expected to reduce or eliminate its activity. Hydrophobic residues are nonpolar amino acids such as phenylalanine, leucine, isoleucine, valine, alanine, methionine, tryptophan, and cysteine. Hydrophilic residues are polar amino acids such as lysine, arginine, histidine, glutamate, and aspartate.

Modifications to a RGS-GEF polypeptide of the present invention or corresponding nucleotide sequence, e.g., mutations, can also be prepared based on homology searching from gene data banks, e.g., Genbank, EMBL. Sequence homology searching can be accomplished using various methods, including algorithms described in the BLAST family of computer programs, the Smith-Waterman algorithm, etc. For example, conserved amino acids can be identified between various sequences containing an RGS domain of various GEF proteins. (See Fig. 18) A mutation(s) can then be introduced into such sequences by identifying and aligning amino acids conserved between the polypeptides and then modifying an amino acid in a conserved or non-conserved position. A mutated RGS-GEF sequence can comprise conserved or non-conserved amino acids, e.g., between corresponding regions of homologous nucleic acids. For example, a mutated sequence can comprise conserved or non-conserved residues from any number of homologous sequences as mentioned-above and/or determined from an appropriate searching algorithm.

Corresponding mutations can be made in specific regions of an RGS-GEF nucleic acid. For example, mutations may be made wherein amino acids that participate in the GTPase catalytic function or mutations may be made in amino acids that function as contact points between the RGS-GEF sequence and the $G\alpha$ subunit.

An RGS-GEF polypeptide or fragment thereof, or substituted RGS-GEF polypeptide or fragment thereof, may also comprise various modifications, wherein such modifications include glycosylation, covalent modifications (e.g., of an R-group of an amino acid), amino acid substitution, amino acid deletion, or amino acid addition. Modifications to the polypeptide can be accomplished according to various methods, including recombinant, synthetic, chemical, etc.

Polypeptides of the present invention (e.g., RGS-GEF polypeptides, and fragments and mutations thereof) may be used in various ways, e.g., as immunogens for antibodies as described below, as biologically-active agents (e.g., having one or more of the activities associated with an RGS-GEF polypeptide), as inhibitors of the activities of the corresponding full-length polypeptide. For example, upon binding of p115 Rho-GEF to the $G\alpha$ subunit, a cascade of events is initiated in the cell, e.g., promoting cell proliferation and/or cytoskeletal rearrangements. The interaction between p115 Rho-GEF and the $G\alpha$ subunit can be modulated by using a RGS-GEF polypeptide, or fragment thereof, to inhibit the interaction between p115 Rho-GEF and the $G\alpha$ subunit. Such a fragment can be useful for modulating pathological conditions associated with the Rho signaling pathway. A useful fragment may be identified routinely by testing the ability of overlapping fragments of the entire length of the RGS domain of a GEF protein to inhibit the binding of p115 Rho-GEF with the $G\alpha$ subunit or to inhibit the GAP activity of the p115 Rho-GEF toward the $G\alpha$ subunit. The measurement of these activities is described below and in the examples. Peptides can be chemically-modified, etc.

A RGS-GEF polypeptide of the present invention can comprise one or more structural domains, functional domains, detectable domains, antigenic domains, and/or other polypeptides of interest, in an arrangement which does not occur in nature, i.e., not naturally-occurring. A polypeptide comprising such features is a chimeric or fusion polypeptide. Such a chimeric polypeptide can be prepared according to various methods, including, chemical, synthetic, quasi-synthetic, and/or recombinant methods. A chimeric nucleic acid coding for a chimeric polypeptide can contain the various domains or desired polypeptides in a continuous or interrupted open reading frame, e.g., containing introns, splice sites, enhancers, etc. The chimeric nucleic acid can be produced according to various methods. See, e.g., U.S. Pat. No. 5,439,819. A domain or desired polypeptide can possess

any desired property, including, a biological function such as catalytic, signaling, growth promoting, cellular targeting, etc., a structural function such as hydrophobic, hydrophilic, membrane-spanning, etc., receptor-ligand functions, and/or detectable functions, e.g., combined with enzyme, fluorescent polypeptide, green fluorescent protein GFP (Chalfie et al., 1994, *Science*, 263:802; Cheng et al., 1996, *Nature Biotechnology*, 14:606; Levy et al., 1996, *Nature Biotechnology*, 14:610, etc. In addition, a RGS-GEF nucleic acid, or a fragment thereof, may be used as selectable marker when introduced into a host cell. For example, a nucleic acid coding for an amino acid sequence according to the present invention can be fused in frame to a desired coding sequence and act as a tag for purification, selection, or marking purposes. The region of fusion encodes a cleavage site.

A polypeptide according to the present invention can be produced in an expression system, e.g., in vivo, in vitro, cell-free, recombinant, cell fusion, etc., according to the present invention. Modifications to the polypeptide imparted by such system include, glycosylation, amino acid substitution (e.g., by differing codon usage), polypeptide processing such as digestion, cleavage, endopeptidase or exopeptidase activity, attachment of chemical moieties, including lipids, phosphates, etc. For example, some cell lines can remove the terminal methionine from an expressed polypeptide.

A polypeptide according to the present invention can be recovered from natural sources, transformed host cells (culture medium or cells) according to the usual methods, including, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, hydroxyapatite chromatography and lectin chromatography. It may be useful to have low concentrations (approximately 0.1-5 mM) of calcium ion present during purification (Price, et al., *J. Biol. Chem.*, 244:917 (1969)). High performance liquid chromatography (HPLC) can be employed for final purification steps.

A RGS-GEF nucleic acid of the present invention can comprise the complete coding sequence for an RGS-GEF polypeptide, or fragments thereof. A nucleic acid according to the present invention may also comprise a nucleotide sequence which is 100% complementary, e.g., an anti-sense, to any RGS-GEF nucleotide sequence.

A nucleic acid according to the present invention can be obtained from a variety of different sources. It may be obtained from DNA or RNA, such as polyadenylated mRNA,

e.g., isolated from tissues, cells, or whole organism. The nucleic acid may be obtained directly from DNA or RNA, or from a cDNA library. The nucleic acid can be obtained from a cell at a particular stage of development, having a desired genotype, phenotype (e.g., an oncogenically transformed cell or a cancerous cell), etc. The nucleic acid may also be
5 chemically synthesized.

A nucleic acid according to the present invention may include only coding sequence for an RGS-GEF polypeptide; coding sequence for an RGS-GEF polypeptide and additional functional coding sequences including, for example, leader sequences, secretory sequences, tag sequences (e.g. targeting tags, enzymatic tags, fluorescent tags etc.). A nucleic acid
10 according to the present invention may also include coding sequence for an RGS-GEF polypeptide and non-coding sequences, e.g., untranslated sequences at either a 5' or 3' end, or dispersed in the coding sequence, e.g., introns.

A nucleic acid according to the present invention may also comprise an expression control sequence operably linked to a nucleic acid as described above. The phrase
15 "expression control sequence" means a nucleic acid sequence which regulates expression of a polypeptide coded for by a nucleic acid to which it is operably linked. Expression can be regulated at the level of the mRNA or polypeptide. Thus, the expression control sequence includes mRNA-related elements and protein-related elements. Such elements include promoters, enhancers (viral or cellular), ribosome binding sequences, transcriptional
20 terminators, etc. An expression control sequence is operably linked to a nucleotide coding sequence when the expression control sequence is positioned in such a manner to effect or achieve expression of the coding sequence. For example, when a promoter is operably linked 5' to a coding sequence, expression of the coding sequence is driven by the promoter. Expression control sequences can be heterologous or endogenous to the normal gene.

25 A nucleic acid in accordance with the present invention may be selected on the basis of nucleic acid hybridization. The ability of two single-stranded nucleic acid preparations to hybridize together is a measure of their nucleotide sequence complementarity, e.g., base-pairing between nucleotides, such as A-T, G-C, etc. The invention thus also relates to nucleic acids which hybridize to a nucleic acids comprising a nucleotide sequence as set
30 forth in Figures 11, 13, 15, and 17. The present invention includes both strands of nucleic acid, e.g., a sense strand and an anti-sense strand.

According to the present invention, a nucleic acid or polypeptide can comprise one or more differences in the nucleotide or amino acid sequence set forth in Figures 10-17. Changes or modifications to the nucleotide and/or amino acid sequence can be accomplished by any method available, including directed or random mutagenesis.

5 A nucleic acid coding for an RGS-GEF polypeptide according to the invention may comprise nucleotides which occur in a naturally-occurring GEF gene e.g., naturally-occurring polymorphisms, normal or mutant alleles (nucleotide or amino acid), mutations which are discovered in a natural population of mammals, such as humans, monkeys, pigs, mice, rats, or rabbits. By the term naturally-occurring, it is meant that the nucleic acid is
10 obtained from a natural source, e.g., animal tissues and cells, body fluids, tissue culture cells, forensic samples. Naturally-occurring mutations include deletions, substitutions, or additions of nucleotide sequence. These genes can be detected and isolated by nucleic acid hybridization according to methods well known to one skilled in the art. It is recognized that, by analogy to other oncogenes, naturally-occurring variants of GEF proteins will
15 include variants with deletions, substitutions, and additions in the RGS domain of a GEF protein, which produce pathological conditions in the host cell and organism.

A nucleotide sequence coding for an RGS-GEF polypeptide of the invention may contain codons found in a naturally-occurring gene, transcript, or cDNA, for example, or it may contain degenerate codons coding for the same amino acid sequences.

20 In addition, a nucleic acid or polypeptide of the present invention may be obtained from any desired mammalian organism, but also non-mammalian organisms. Homologs from mammalian and non-mammalian organisms can be obtained according to various methods. For example, hybridization with an oligonucleotide (see below) selective for an RGS domain of a GEF, or a RGS-GEF, of the present invention can be employed to select
25 such homologs, e.g., as described in Sambrook et al., *Molecular Cloning*, 1989, Chapter 11. Such homologs may have varying amounts of nucleotide and amino acid sequence identity and similarity to previously identified RGS domain or RGS-GEF nucleotide or polypeptide sequence. Non-mammalian organisms include, e.g., vertebrates, invertebrates, zebra fish, chicken, *Drosophila*, yeasts (such as *Saccharomyces cerevisiae*), *C. elegans*, roundworms,
30 prokaryotes, plants, *Arabidopsis*, viruses, etc.

A nucleic acid according to the present invention may comprise, for example, DNA, RNA, synthetic nucleic acid, peptide nucleic acid, modified nucleotides, or mixtures thereof.

A DNA can be double- or single-stranded. Nucleotides comprising a nucleic acid can be joined via various known linkages such as, for example, ester, sulfamate, sulfamide, phosphorothioate, phosphoramidate, methylphosphonate, carbamate, etc., depending on the desired purpose. Linkages may be modified for purposes such as, for example, resistance to nucleases such as RNase H and improved *in vivo* stability. See, e.g., U.S. Pat. Nos. 5,378,825.

Various modifications can be made to the nucleic acids, such as attaching detectable markers (avidin, biotin, radioactive elements), moieties which improve hybridization, detection, or stability. The nucleic acids can also be attached to solid supports, e.g., nitrocellulose, nylon, agarose, diazotized cellulose, latex solid microspheres, polyacrylamides, etc., according to a desired method. See, e.g., U.S. Pat. Nos. 5,470,967, 5,476,925, 5,478,893.

Another aspect of the present invention relates to oligonucleotides and nucleic acid probes. Such oligonucleotides or nucleic acid probes can be used, e.g., to detect, quantify, or isolate an RGS-GEF nucleic acid in a test sample. Detection can be desirable for a variety of different purposes, including research, diagnostic, and forensic. For diagnostic purposes, it may be desirable to identify the presence or quantity of a specific RGS-GEF nucleic acid sequence in a sample obtained from tissues, cells, body fluids, etc. In a preferred method, the present invention relates to a method of detecting a target RGS-GEF nucleic acid in a test sample comprising contacting the test sample with an oligonucleotide under conditions effective to achieve hybridization between the target and oligonucleotide; and detecting hybridization. An oligonucleotide in accordance with the invention can also be used in synthetic nucleic acid amplification such as PCR, e.g., Saiki et al., 1988, *Science*, 241:53; U.S. Pat. No. 4,683,20, or or differential display (See, e.g., Liang et al., *Nucl. Acid. Res.*, 21:3269-3275, 1993; USP 5,599,672; WO97/18454). Oligonucleotides can be identified routinely, e.g., to the DH, PH, and RGS-GEF domains to differentially display and/or amplify gene products containing such sequences.

Both sense and antisense nucleotide sequences are intended as part of the invention. A unique nucleic acid according to the present invention may be determined routinely. An

RGS-GEF nucleic acid may be used as a hybridization probe to identify the presence of RGS-GEF nucleotide sequence in a sample comprising a mixture of nucleic acids, e.g., on a Northern blot. Hybridization can be performed under stringent conditions to select nucleic acids having at least 95% identity (i.e., complementarity) to the probe, but less stringent
5 conditions can also be used. A unique RGS-GEF nucleotide sequence can also be fused in-frame, at either its 5' or 3' end, to various nucleotide sequences, including, for example, coding sequences for enzymes or expression control sequences, etc.

Hybridization can be performed under different conditions, depending on the desired selectivity, e.g., as described in Sambrook et al., *Molecular Cloning*, 1989. For example, to
10 specifically detect RGS-GEF sequences, an oligonucleotide can be hybridized to a target nucleic acid under conditions in which the oligonucleotide only hybridizes to the GEF sequence from which the RGS-GEF sequence was derived, e.g., where the oligonucleotide is 100% complementary to the target. Different conditions can be used if it is desired to select target nucleic acids which have less than 100% nucleotide complementarity, at least
15 about, e.g., 99%, 97%, 95%, 90%, 70%, 67%. Since a mutation in GEF genes can cause diseases or pathological conditions, e.g., cancer, benign tumors, an oligonucleotide according to the present invention can be used diagnostically. For example, a patient having symptoms of a cancer or other condition associated with the Rho signaling pathway (see below) can be diagnosed with the disease by using an oligonucleotide according to the
20 present invention, in polymerase chain reaction followed by DNA sequencing to identify whether the sequence is normal, in combination with other oncogene oligonucleotides, etc., e.g., p53, Rb, p21, Dbl, MTS1, Wt1, Bcl-1, Bcl-2, MDM2, etc.

Oligonucleotides according to the present invention can be of any desired size, preferably 14-16 oligonucleotides in length, or more. Such oligonucleotides can have non-
25 naturally-occurring nucleotides, e.g., inosine. In accordance with the present invention, the oligonucleotide can comprise a kit, where the kit includes a desired buffer (e.g., phosphate, tris, etc.), detection compositions, etc. The oligonucleotide can be labeled or unlabeled, with radioactive or non-radioactive labels as known in the art.

Anti-sense nucleic acid can also be prepared from a nucleic acid according to the
30 present, preferably an anti-sense RGS-GEF nucleotide sequence corresponding to an RGS-GEF nucleotide sequence of Figures 11, 13, 15, and 17. Anti-sense RGS-GEF nucleic acid

can be used in various ways, such as to regulate or modulate expression of GEF proteins containing RGS domains or to detect expression of RGS-GEF proteins, including by *in situ* hybridization. For the purposes of regulating or modulating expression, an anti-sense oligonucleotide may be operably linked to an expression control sequence.

5 The RGS-GEF nucleic acids according to the present invention can be labelled according to any desired method. The nucleic acid can be labeled using radioactive tracers such as ^{32}P , ^{35}S , ^{125}I , ^3H , or ^{14}C , to mention only the most commonly used tracers. The radioactive labeling can be carried out according to any method such as, for example, terminal labeling at the 3' or 5' end using a radiolabeled nucleotide, polynucleotide kinase
10 (with or without dephosphorylation with a phosphatase) or a ligase (depending on the end to be labeled). A non-radioactive labeling can also be used, combining a nucleic acid of the present invention with residues having immunological properties (antigens, haptens), a specific affinity for certain reagents (ligands), properties enabling detectable enzyme reactions to be completed (enzymes or coenzymes, enzyme substrates, or other substances
15 involved in an enzymatic reaction), or characteristic physical properties, such as fluorescence or the emission or absorption of light at a desired wavelength, etc.

 An RGS-GEF nucleic acid according to the present invention, including oligonucleotides, anti-sense nucleic acid, etc., can be used to detect expression of RGS-GEF nucleic acids in whole organs, tissues, cells, etc., by various techniques, including Northern
20 blot, PCR, *in situ* hybridization, etc. Such nucleic acids can be particularly useful to detect disturbed expression, e.g., cell-specific and/or subcellular alterations of RGS-GEF expression. The levels of RGS-GEF proteins can be determined alone or in combination with other genes products (oncogenes such as p53, Rb, Wt1, etc.), transcripts, etc.

 A nucleic acid according to the present invention can be expressed in a variety of
25 different systems, *in vitro* and *in vivo*, according to the desired purpose. For example, a nucleic acid can be inserted into an expression vector, introduced into a desired host, and cultured under conditions effective to achieve expression of a polypeptide coded for the nucleic acid. Effective conditions includes any culture conditions which are suitable for achieving production of the polypeptide by the host cell, including effective temperatures,
30 pH, medias, additives to the media in which the host cell is cultured (e.g., additives which amplify or induce expression such as butyrate, or methotrexate if the coding nucleic acid is

adjacent to a dhfr gene), cyclohexamide, cell densities, culture dishes, etc. A nucleic acid can be introduced into the cell by any effective method including, e.g., calcium phosphate precipitation, electroporation, injection, DEAE-Dextran mediated transfection, fusion with liposomes, and viral transfection. A cell into which a nucleic acid of the present invention
5 has been introduced is a transformed host cell. The nucleic acid can be extrachromosomal or integrated into a chromosome(s) of the host cell. It can be stable or transient. An expression vector is selected for its compatibility with the host cell. Host cells include, mammalian cells, e.g., COS-7, CHO, HeLa, LTK, NIH 3T3, yeast, insect cells, such as Sf9 (*S. frugipeda*) and *Drosophila*, bacteria, such as *E. coli*, *Streptococcus sp.*, *Bacillus sp.*,
10 yeast, fungal cells, plants, embryonic stem cells (e.g., mammalian, such as mouse or human), cancer or tumor cells. Sf9 expression can be accomplished in analogy to Graziani et al., *Oncogene*, 7:229-235, 1992. Expression control sequences are similarly selected for host compatibility and a desired purpose, e.g., high copy number, high amounts, induction, amplification, controlled expression. Other sequences which can be employed include
15 enhancers such as from SV40, CMV, inducible promoters, cell-type specific elements, or sequences which allow selective or specific cell expression.

A labelled polypeptide can be used, e.g., in binding assays, such as to identify substances that bind or attach to p115 Rho-GEF, to track the movement of p115 Rho-GEF in a cell, in an *in vitro*, *in vivo*, or *in situ* system, etc.

20 A nucleic acid or polypeptide of the present invention can also be substantially purified. By substantially purified, it is meant that nucleic acid or polypeptide is separated and is essentially free from other nucleic acids or polypeptides, i.e., the nucleic acid or polypeptide is the primary and active constituent.

Another aspect of the present invention relates to the regulation of biological
25 pathways in which a RGS-GEF polypeptide is involved, particularly pathological conditions, e.g., cell proliferation (e.g., cancer), growth control, morphogenesis, stress fiber formation, and integrin-mediated interactions, such as embryonic development, tumor cell growth and metastasis, programmed cell death, hemostasis, leucocyte homing and activation, bone resorption, clot retraction, and the response of cells to mechanical stress. See, e.g., Clark
30 and Brugge, *Science*, 268:233-239, 1995; Bussey, *Science*, 272:225-226, 1996. Thus, the invention relates to all aspects of a method of modulating an activity of a RGS-GEF

polypeptide comprising, administering an effective amount of an RGS-GEF polypeptide or a biologically-active fragment thereof, an effective amount of a compound which modulates the activity of a RGS-GEF polypeptide, or an effective amount of a nucleic acid which codes for a RGS-GEF polypeptide or a biologically-active fragment thereof. The activity of the RGS-GEF which is modulated may include binding to a G α subunit or GAP activity toward a G α subunit. The activity can be modulated by increasing, reducing, antagonizing, or promoting expression or activity of the RGS-GEF.

The present invention also relates to antibodies which specifically recognize a RGS-GEF polypeptide. Antibodies, e.g., polyclonal, monoclonal, recombinant, chimeric, can be prepared according to any desired method. For example, for the production of monoclonal antibodies, an RGS-GEF polypeptide according to Figures 10, 12, 14, or 16 can be administered to mice, goats, or rabbit subcutaneously and/or intraperitoneally, with or without adjuvant, in an amount effective to elicit an immune response. The antibodies can also be single chain or FAb. The antibodies can be IgG, subtypes, IgG2a, IgG1, etc.

An antibody specific for RGS-GEF means that the antibody recognizes a defined sequence of amino acids within or including the amino acid sequence of the RGS domain of a GEF polypeptide. Thus, a specific antibody will bind with higher affinity to an amino acid sequence, i.e., an epitope, found in the RGS domain of a GEF polypeptide than to a different epitope(s), e.g., as detected and/or measured by an immunoblot assay. Thus, an antibody which is specific for an epitope within or including the RGS domain of p115 Rho-GEF is useful to detect the presence of the epitope in a sample, e.g., a sample of tissue containing p115 Rho-GEF gene product, distinguishing it from samples in which the epitope is absent.

Additionally, in accordance with the present invention, ligands which bind to an RGS domain of a GEF polypeptide can also be prepared, e.g., using synthetic peptide libraries or aptamers (e.g., Pitrun et al., U.S. Pat. No. 5,143,854; Geysen et al., 1987, J. Immunol. Methods, 102:259-274; Scott et al., 1990, Science, 249:386; Blackwell et al., 1990, Science, 250:1104; Tuerk et al., 1990, Science, 249: 505).

Antibodies and other ligands which bind the RGS domain of a GEF polypeptide, and specifically antibodies and other ligands which bind the RGS domain of p115 Rho GEF, can be used in various ways. These include, but are not limited to, uses therapeutic, diagnostic, and commercial research tools, e.g, to quantitate the levels of p115 Rho-GEF polypeptide in

animals, tissues, cells, etc., to identify the cellular localization and/or distribution of p115 Rho-GEF, to purify p115 Rho-GEF or a polypeptide comprising a part of p115 Rho-GEF, to modulate the function of p115 Rho-GEF, etc. Antibodies can be used in Western blots, ELIZA, immunoprecipitation, RIA, etc. The present invention relates to such assays,
5 compositions and kits for performing them, etc.

An antibody according to the present invention can be used to detect polypeptides or fragments containing an RGS domain of a GEF polypeptide in various samples, including tissue, cells, body fluid, blood, urine, cerebrospinal fluid. A method of the present invention comprises contacting a ligand which binds to an RGS-GEF polypeptide of Figure 10, 12, 14,
10 or 16 under conditions effective, as known in the art, to achieve binding, detecting specific binding between the ligand and peptide. By specific binding, it is meant that the ligand attaches to a defined sequence of amino acids, e.g., within or including the amino acid sequence of the RGS domain as shown in Figures 10, 12, 14, and 16, or derivatives thereof. The antibodies or derivatives thereof can also be used to inhibit expression of GEF proteins
15 containing an RGS domain. The levels of a GEF polypeptide containing an RGS domain may be determined alone or in combination with other gene products. In particular, the amount (e.g., its expression level) of the GEF polypeptide containing an RGS domain can be compared (e.g., as a ratio) to the amounts of other polypeptides in the same or different sample, e.g., p21, p53, Rb, WT1, etc.

20 A ligand for the RGS domain of GEF polypeptides can be used in combination with other antibodies, e.g., antibodies that recognize oncological markers of cancer, including, Rb, p53, c-erbB-2, oncogene products, etc. In general, reagents which are specific for the RGS domain of GEF polypeptides can be used in diagnostic and/or forensic studies according to any desired method, e.g., as U.S. Pat. Nos. 5,397,712; 5,434,050; 5,429,947.

25 The present invention also relates to a transgenic animal, e.g., a non-human-mammal, such as a mouse, comprising an RGS-GEF polypeptide. Transgenic animals can be prepared according to known methods, including, e.g., by pronuclear injection of recombinant genes into pronuclei of 1-cell embryos, incorporating an artificial yeast chromosome into embryonic stem cells, gene targeting methods, embryonic stem cell
30 methodology. See, e.g., U.S. Patent Nos. 4,736,866; 4,873,191; 4,873,316; 5,082,779; 5,304,489; 5,174,986; 5,175,384; 5,175,385; 5,221,778; Gordon et al., *Proc. Natl. Acad.*

5 *Sci.*, 77:7380-7384 (1980); Palmiter et al., *Cell*, 41:343-345 (1985); Palmiter et al., *Ann. Rev. Genet.*, 20:465-499 (1986); Askew et al., *Mol. Cell. Bio.*, 13:4115-4124, 1993; Games et al. *Nature*, 373:523-527, 1995; Valancius and Smithies, *Mol. Cell. Bio.*, 11:1402-1408, 1991; Stacey et al., *Mol. Cell. Bio.*, 14:1009-1016, 1994; Hasty et al., *Nature*, 350:243-246, 1995; Rubinstein et al., *Nucl. Acid Res.*, 21:2613-2617, 1993. A nucleic acid according to the present invention can be introduced into any non-human mammal, including a mouse (Hogan et al., 1986, in *Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York), pig (Hammer et al., *Nature*, 315:343-345, 1985), sheep (Hammer et al., *Nature*, 315:343-345, 1985), cattle, rat, or
10 primate. See also, e.g., Church, 1987, *Trends in Biotech.* 5:13-19; Clark et al., 1987, *Trends in Biotech.* 5:20-24, and DePamphilis et al., 1988, *BioTechniques*, 6:662-680. Additionally, custom transgenic rat and mouse production is commercially available. These transgenic animals are useful, for example, as a cancer model or as a model to evaluate the effects of overexpression of the RGS-GEF polypeptide.

15 Generally, the nucleic acids, polypeptides, antibodies, etc. of the present invention can be prepared and used as described in, U.S. Pat. Nos. 5,501,969, 5,506,133, 5,441,870; WO 90/00607; WO 91/15582;

Other aspects of this invention relate to methods to assay for, or identify, molecules that modulate the following interactions and effects: the interaction between an RGS domain
20 of a GEF and its cognate binding substrate; the interaction between an RGS domain of a GEF and a G α subunit; the effect of G protein subunit stimulation on a guanine nucleotide exchange activity of a GEF protein containing an RGS domain; the effect of a GEF protein having an RGS domain as a GTPase activating protein for a G protein subunit.

Activity can be modulated in various ways, e.g., enhancing, activating, stimulating,
25 suppressing, preventing, inhibiting, etc. A modulatory molecule can be an agonist, antagonist, or have partial activities thereof. Modulating molecules can be any type of molecule, including but not limited to small molecules, proteins, peptides, antibodies, nucleic acids, etc. In general, a compound having an *in vitro* activity will be useful *in vivo* to modulate a biological pathway associated with a GEF protein containing an RGS domain,
30 e.g., to treat a pathological condition associated with the biological and cellular activities

mentioned above. The modulatory molecules can comprise a mixture of the same or different molecules.

A binding substrate for the RGS domain of a GEF protein can be any material to which the RGS domain binds specifically, including members of the $G\alpha_{12}$ family. See, 5 e.g., Strathman and Simon, *Proc. Natl. Acad. Sci.*, 88:5582, 1991. For example, a method of identifying or assaying for a molecule that modulates or regulates the binding of a G protein α subunit to a GEF protein containing an RGS domain, such as p115 Rho-GEF, can be conducted in accordance with this invention. In one embodiment, a GTP bound α subunit, or derivative thereof, is incubated with a GEF protein, or fragment thereof, 10 containing the RGS domain, in the presence and absence of a test molecule to determine whether the presence of the test compound modulates the binding between the GEF protein and the G protein α subunit. The incubation is accomplished under effective conditions, i.e., conditions under which binding or attachment occurs. Binding can be detected in one or more ways. For example, the GEF protein or the binding substrate is labeled detectably; 15 the labelled bound component is separated from the labelled free component; and the amount of bound-detectably labeled GEF protein or binding substrate determined. The detectable label can be of any desired composition, e.g., radioactive, fluorescent, etc. Such an assay can be performed in either solid or liquid phase.

In one aspect of the invention, it is desirable to identify molecules that regulate the 20 binding of the $G\alpha_{12}$ family of subunits, eg. $G\alpha_{12}$ and $G\alpha_{13}$, with a GEF, e.g., p115 Rho GEF, Lsc, KIAA380, or DrhoGEF2. The assay can be conducted using a complete GEF protein, or any subfragments thereof which contain the RGS domain, or biologically active subfragments of the RGS domain. The assay is typically conducted with stable analogs of the GTP bound state of the $G\alpha$ subunit, including α subunits bound to either $GDP\text{-}AlF_4^-$ or 25 $GTP\gamma S$. For example, a binding assay may be conducted by the procedure described in Example 5 below wherein a COS cell is transfected with a nucleic acid construct for a myc-tagged polypeptide, such as p115 Rho GEF, or a fragment thereof and complexes of the polypeptide and a $G\alpha$ subunit are detected by precipitation of any bound complex with a first antibody to one of components and detection of the amount of a second bound component 30 with a second antibody. Binding assays could also be performed using techniques that are

well known in the art such as by binding one of the components to a column and then determining the amount of a second labelled component that binds to the column. Relevant assay methods are also disclosed, for example, in Berman et al, 1996, *J. Biol. Chem.* 271:27209.

5 A method of isolating or assaying for a molecule that modulates or regulates the stimulatory effect of a RGS-GEF polypeptide on GTPase activity, such as a GTPase activity of a $G\alpha$ subunit, can also be conducted in accordance with the invention. For example, a $G\alpha$ subunit is incubated under effective conditions with an RGS-GEF polypeptide having GTPase stimulatory effect in the presence and absence of a test inhibitor to determine
10 whether the presence of the test inhibitor modulates its stimulatory effect. The assay can be conducted using a complete RGS-GEF polypeptide, a GEF protein, or any subfragments thereof which contain the RGS domain, or biologically active subfragments of the RGS domain. An RGS-GEF polypeptide can be p115 Rho GEF, Lsc, KIAA380, DrhoGEF2, or biologically-active fragments thereof. For example, an assay can be conducted using a p115
15 Rho GEF in conjunction with an $\alpha 12$ or an $\alpha 13$ subunit, as described in the examples discussed herein, as well as using other variations or assay methods which are well known in the art. For example, the assay may be conducted in accordance with Example 5 below, in which $G\alpha$ subunits were loaded with $[\gamma\text{-}^{32}\text{P}]\text{GTP}$ and the amount of hydrolysis under various conditions, including the presence of an RGS-GEF polypeptide, was determined by
20 measuring the amount of ^{32}Pi in the supernatant after centrifugation of the assay mixture. Relevant assay methods are also disclosed, for example, in Berman et al, 1996, *J. Biol. Chem.* 271:27209.

 A method of identifying or assaying for a molecule that modulates the stimulatory effect of an activated $G\alpha$ subunit on a RGS-GEF polypeptide having GEF mediated
25 nucleotide exchange for a monomeric G protein can also be conducted in accordance with this invention. For instance, a first assay can be conducted by incubating an activated $G\alpha$ alpha subunit with a GEF protein (e.g., p115 Rho GEF, Lsc, KIAA380, DrhoGEF2, or biologically-active fragments thereof, which retain GEF activity) and a monomeric G protein in the presence and absence of a test modulator to determine whether the test
30 modulator has an inhibitory, enhancing, etc. effect on the ability of an activated $G\alpha$ subunit

to stimulate GEF mediated nucleotide exchange of a monomeric protein. See e.g. Hart et al., 1996, *J. Biol. Chem.* 221:25452. The test modulator can be further evaluated by conducting a second assay in which said GEF protein and a monomeric G protein, without the G protein subunit, are incubated in the presence or absence of the test modulator to determine whether the test modulator had any effect on GEF mediated nucleotide exchange of the monomeric protein, and then comparing the modulation effect in the first and second assays to determine whether the modulating effect in the first assay is different from the modulating effect in the second assay, thereby indicating that the test modulator modulates the interaction of an activated $G\alpha$ subunit with the GEF protein rather than the interaction of the GEF protein with the monomeric G protein. For example, the stimulatory effect on GEF mediated guanine nucleotide exchange may be measured according to Example 6 below, wherein RhoA was loaded with [3H]GDP and the dissociation of GDP from RhoA was measured under various conditions by the determination of bound GDP by filtration, prior to an after incubation. (See e.g. Northrup et al., *J. Biol. Chem.*, 257, 11416-11423 (1982)).

A method of identifying a molecule that mimics the stimulatory effect of an activated $G\alpha$ subunit on GEF mediated nucleotide exchange of a monomeric G protein may also be conducted in accordance with the invention. The method comprises identifying a test compound that exhibits a binding affinity for the RGS domain of GEF proteins and then incubating a GEF protein and monomeric G protein in the presence or absence of the test compound to determine whether the test compounds exhibits a stimulatory effect on GEF mediated nucleotide exchange of a monomeric G protein. The identification of test compounds that exhibit a binding affinity for the RGS domain of GEF proteins may be accomplished using techniques well known in the art. For example, an RGS polypeptide may be bound to a column and cocktails of test compounds may be passed over the column to determine if any were selectively bound by the column.

A method of identifying a molecule, or mixture of molecules, that mimics the stimulatory effect of an RGS domain of GEF polypeptide on $G\alpha$ subunit GTPase activity may also be conducted in accordance with the invention. The method comprises identifying a test compound that exhibits a binding affinity for a $G\alpha$ subunit and incubating a GTP loaded $G\alpha$ subunit in the presence or absence of the test compound to determine whether the test compound exhibits a stimulatory effect GTPase activity of the $G\alpha$ subunit. The

identification of test compounds that exhibit a binding affinity for the G α subunit may be accomplished using techniques well known in the art. For example, a G α_{12} may be bound to a substrate and incubated with both a GEF polypeptide containing an RGS domain and the test compound to determine whether the test compound competes with the RGS domain for binding to the G α subunit.

The modulation of oncogenic transforming activity by an RGS-GEF component, or derivatives thereof, can be measured according to various known procedures, e.g., Eva and Aaronson, *Nature*, 316:273-275, 1985; Hart et al., *J. Biol. Chem.*, 269:62-65, 1994. A compound can be added at any time during the method (e.g., pretreatment of cells; after addition of the RGS-GEF, etc.) to determine its effect on the oncogenic transforming activity of the RGS-GEF component. Various cell lines can also be used.

Other assays for monomeric GTPase-mediated signal transduction can be accomplished according to the invention by analogy to procedures known in the art, e.g., as described in U.S. Pat. Nos. 5,141,851; 5,420,334; 5,436,128; and 5,482,954; WO94/16069; WO93/16179; WO91/15582; WO90/00607.

The present invention thus also relates to the treatment and prevention of diseases and pathological conditions associated with signal transduction mediated by GEF proteins that contain an RGS domain, e.g., cancer, diseases associated with abnormal cell proliferation. For example, the invention relates to a method of treating cancer comprising administering, to a subject in need of treatment, an amount of a compound effective to treat the disease, where the compound is a regulator of the stimulatory effect of GEF protein containing an RGS on G α subunit GTPase activity or where the compound is a regulator of the stimulatory effect of a G α subunit on GEF mediated nucleotide exchange by a monomeric GTPase. Treating the disease can mean, delaying its onset, delaying the progression of the disease, improving or delaying clinical and pathological signs of disease. A regulator compound, or mixture of compounds, can be synthetic, naturally-occurring, or a combination. A regulator compound can comprise amino acids, nucleotides, hydrocarbons, lipids, polysaccharides, etc. A regulator compound is preferably a compound that regulates expression of a GEF protein containing an RGS domain, e.g., inhibiting or increasing its mRNA, protein expression, or processing, or a compound that regulates the interaction of the RGS domain of the GEF protein with a G α subunit. To treat the disease, the compound,

or mixture, can be formulated into pharmaceutical composition comprising a pharmaceutically acceptable carrier and other excipients as apparent to the skilled worker. See, e.g., *Remington's Pharmaceutical Sciences*, Eighteenth Edition, Mack Publishing Company, 1990. Such composition can additionally contain effective amounts of other
5 compounds, especially for treatment of cancer.

EXAMPLES

Example 1. Identification of homology between a Rho GEF and proteins which regulate G-protein signaling.

The RGS family of proteins act as negative regulators of G protein signalling.
10 Nineteen mammalian members of the family have been identified, all of which encode proteins that contain a homologous core domain called the RGS box.

Examination of the sequence of p115-GEF, a GEF specific for Rho, revealed an N-terminal region with specific homology to the conserved domain of RGS proteins, including RGS4, RGS2, GAIP, RGS12, and RGS14 (Fig. 1). Analysis of three other Rho GEF
15 proteins, Lsc, KIAA380, and DrhoGEF also showed that they contained regions of specific homology to the conserved domain of RGS proteins (Fig. 1).

The crystal structure of a complex between RGS4 and AlF_4 -activated $\text{G}\alpha_i$ revealed that the functional core of RGS4 (the RGS box) contains nine α -helices that fold into two small subdomains (Tesmer et al., *Cell*, **89**, 251 (1997)). The RGS box has been shown to
20 contain the GAP activity towards $\text{G}\alpha$ subunits (Popov et al., *Proc. Natl. Acad. Sci. USA*, **94**, 7216 (1997)). The hydrophobic core residues of the box, which are conserved in members of the RGS family, are important for stability of structure and GAP activity (Tesmer et al., *Cell*, **89**, 251 (1997) and Srinivasan et al., *J. Biol. Chem.*, **273**, 1529 (1998). RGS4 stimulates the GTPase activity of $\text{G}\alpha_i$ by interacting with its three switch regions, primarily
25 by stabilization of the transition state of GTP hydrolysis (Tesmer et al., *Cell*, **89**, 251 (1997)).

Most of the hydrophobic residues that form the core of the RGS domain are conserved in p115 Rho GEF (17 out of 23) (Fig. 1). The position of gaps in the alignment correspond to the loops between alpha helices of RGS domain structure. This homology
30 suggested that the N-terminal region of p115-GEF may have a similar structure to the RGS4 box domain and possess GAP activity. In contrast, the residues of RGS4 that make contact

with the switch regions of $G\alpha_{i1}$ (GDP- AlF_4^-) are not well conserved, and any GAP activity of p115 Rho GEF will have a unique mechanism or a significantly different specificity than those previously identified.

A search of the gene bank revealed three other Rho-GEF members that have regions homologous to the RGS region of p115. These include Lsc, KIAA380, and DrhoGEF2 (Fig. 1). Lsc appears to be the mouse homologue of p115 Rho GEF and KIAA380 appears to be the human homologue of *Drosophila* DrhoGEF2 (Whitehead et al., *J. Biol. Chem.*, **271**, 18643 (1996); Barrett et al., *Cell*, **91**, 905 (1997)). These four Rho-GEF's define a new RGS related family of proteins which also possess guanine nucleotide exchange activity for Rho.

An alignment of RGS domains of the four GEF proteins known to contain RGS domains (p115 Rho GEF, Lsc, KIAA380, DRhoGEF2) with the RGS domains of RGS proteins RET-RGS1, RGS1, RGS2, RGS3, RGS4, RGS7, RGS10, RGS12, RGS14, Rap1/2B.P., and GAIIP shows that a novel sub-RGS consensus sequence is defined by the RGS sequence of the four GEF proteins (Fig. 18). As shown in the bottom set of sequences shown in Fig. 18, a novel sub-RGS consensus sequence is shown by the large gap of 13 to 14 amino acids in the homology alignment, along with the conservation of amino acids on either side of the gap.

Example 2. The RHO GEF protein, p115 RHO-GEF, stimulates the GTPase activity of $G\alpha_{i3}$ and $G\alpha_{i2}$ subunits.

P115 Rho GEF was tested to determine its capability in stimulating the intrinsic GTPase activity (GAP activity) of $G\alpha_{i3}$ and $G\alpha_{i2}$.

$G\alpha_{i2}$ was expressed in Sf9 cells and purified as described in Kozasa and Gilman, *J. Biol. Chem.*, **270**, 1734 (1995). $G\alpha_{i3}$ was prepared by a similar procedure using the previously described baculovirus method (Singer and Miller, *J. Biol. Chem.*, **269**, 19796 (1994)) and octylglucoside during washing and elution of the α subunit after immobilization of the heterotrimer on Ni-NTA resin (Qiagen). The eluted $G\alpha_{i3}$ was further purified by absorption to and elution from hydroxyapatite. $G\alpha_{i2}$ or $G\alpha_{i3}$ (20-30 pmol) was loaded at 30°C for 30 or 40 minutes, respectively with 5 μ M [γ - 32 P]GTP (50-100 cpm/fmol) and in the

presence of 5 mM EDTA. Samples were then rapidly filtered by centrifugation at 4°C through Sephadex G50 which had been equilibrated with buffer A (50 mM NaHepes (pH 8.0), 1 mM dithiothreitol, 5 mM EDTA, and 0.05% polyoxyethylene 10-laurylether) to remove free [γ - 32 P]GTP and [32 Pi]. Hydrolysis of GTP was initiated by adding G α loaded with [γ - 32 P]GTP in buffer A containing 8 mM MgSO₄, 1 mM GTP and the indicated amount of p115. The reaction mixture was incubated at 4°C or 15°C. Aliquots (50 μ l) were removed at the indicated times and mixed with 750 μ l of 5%(w/v) NoritA in 50 mM NaH₂PO₄. The mixture was centrifuged at 2000 rpm for 5 minutes and 400 μ l of supernatant containing 32 Pi were counted by liquid scintillation spectrometry.

The hydrolysis of GTP bound to G α_{13} and G α_{12} was performed at 15°C either with or without 10nM full-length p115 (Fig. 2, Panel A). The hydrolysis of GTP bound G α_{13} and G α_{12} was measured at 4°C in the presence of various concentrations of p115 (Fig. 2, Panel B). Full-length p115 was able to stimulate a single round of hydrolysis of [γ - 32 P]GTP which had been prebound to the G α_{13} subunit. The intrinsic GTPase activity of G α_{12} , the closest homologue of G α_{13} , was also stimulated by full-length p115. At 15°C, the k_{cat} for hydrolysis of GTP by G α_{12} (0.07 min⁻¹) and G α_{13} (0.24 min⁻¹) were respectively increased 5-fold and 10-fold by 10 nM p115 (Fig. 2, Panel A). Similar results were obtained with several preparations of G α_{12} and G α_{13} . Treatment of p115 at 90°C inactivated this GAP activity. Due to the rapid hydrolytic rates of G α_{13} , assays were performed at 4°C to better estimate the effect of p115 on the initial rate of GTPase activity by the G protein (Fig. 2, Panel B). Under these conditions, 100 nM p115 caused an 80-fold increase in the GTPase activity of G α_{13} . In contrast, the hydrolytic rate of G α_{12} was increased only 6-fold. Although stimulation of both proteins was observed at concentrations of p115 as low as 1 nM, measurements at both temperatures indicate that p115 is a more efficacious GAP for G α_{13} than G α_{12} .

In the absence of a receptor, the rate limiting step in the binding of GTP γ S to G α and the steady state hydrolysis of GTP is the release of GDP. P115 did not affect either the rate of GTP γ S binding to G α_{12} and G α_{13} or the steady state of GTPase activity of either subunit.

Therefore, p115 stimulates only the intrinsic GTPase activity of $G\alpha_{12}$ and $G\alpha_{13}$ without effecting their rates of nucleotide exchange.

The conserved RGS box region of RGS proteins is sufficient to show GAP activity *in vitro* (Popov et al., *Proc. Natl. Acad. Sci. USA*, **94**, 7216 (1997)). Therefore, a fusion protein (Fig. 1, Panel B) of glutathione-S-transferase and the N-terminal region of p115, GST-RGS, was tested for GAP activity. This region retains RGS homology domain but not the Dbl or PH domains of p115. This "RGS domain" of p115 (10 nM) was almost as active as full-length p115 when tested for GAP activity for $G\alpha_{12}$ and $G\alpha_{13}$ (Fig. 3). In contrast, a construct of p115 missing this N-terminal region was ineffective. Thus, the data indicates that the RGS homology region is responsible for the GAP activity of p115.

Example 3. The p115 RHO-GEF, does not stimulate the GTPase activity of $G\alpha_i$, $G\alpha_z$, $G\alpha_q$ and $G\alpha_s$ subunits.

The specificity of the GAP activity of p115 for various G protein α subunits was examined as follows.

$G\alpha_s$ was expressed in and purified from *Escherichia coli* as described in Lee et al., *Meth. Enzymol.*, **237**, 146 (1994). $G\alpha_i$, $G\alpha_z$, and $G\alpha_q$ R183C were expressed in Sf9 cells and purified as described in Kozasa and Gilman, *J. Biol. Chem.*, **270**, 1734 (1995) and Biddlecome et al., *J. Biol. Chem.*, **271**, 7999 (1996). $G\alpha_i$, $G\alpha_z$, and $G\alpha_z$ were loaded with 5-10 μ M [γ - 32 P]GTP at 20°C (for $G\alpha_s$) or 30°C (for $G\alpha_i$ and $G\alpha_z$) for 20 minutes in the presence of 5 mM EDTA and GAP assays were performed as described above for $G\alpha_{12}$ and $G\alpha_{13}$. Gap activity on $G\alpha_q$ was assessed with a mutant $G\alpha_q$ R183C. An analogous mutation in $G\alpha_i$ R1178C, causes markedly reduced GTPase activity but response to RGS proteins was retained (Berman et al., *Cell*, **86**, 445 (1996)). The slow GTPase activity of $G\alpha_q$ R183C enables loading of [γ - 32 P]GTP on $G\alpha_q$ without using receptor to accelerate nucleotide exchange. $G\alpha_q$ R183C was loaded with 10 μ M [γ - 32 P]GTP in the presence of 50 mM Hepes (pH7.4), 0.1 mg/ml BSA, 1 mM DTT, 1 mM EDTA, 0.9 mM $MgSO_4$, 30 mM $(NH_4)_2SO_4$, 4% glycerol, and 5.5 mM CHAPS at 20°C for 2 hours. The reaction mixture was rapidly

filtered through Sephadex G50 which had been equilibrated with 50 mM Hepes (pH 7.4), 1 mM DTT, 1 mM EDTA, 0.9 mM SO_4 , 0.1 mg/ml BSA, and 1 mM CHAPS.

The results of this study showed that p115 (100 nM) did not stimulate the GTPase activity of $\text{G}\alpha_i$, $\text{G}\alpha_z$, or $\text{G}\alpha_q$ under conditions where RGS4 acts as a GAP for these $\text{G}\alpha$ subunits (Figure 4). Similarly, p115 did not accelerate the GTPase activity of $\text{G}\alpha_s$, nor did p115 Rho GEF have any GAP activity towards RhoA or rac1. Thus, p115 is a GAP with specificity for $\text{G}\alpha_{12}$ and $\text{G}\alpha_{13}$.

Example 4. Selective inhibition of p115 GAP activity by AlF_4^- activated forms of $\text{G}\alpha$ subunits.

RGS proteins have been shown to have high affinity for the GDP-AlF_4^- bound form of α subunits, a configuration similar to the transition state of GTP hydrolysis (Tesmer et al., *Cell*, **89**, 251 (1997), Berman et al., *J. Biol. Chem.*, **271**, 27209 (1996)). Therefore, the GDP-AlF_4^- forms of $\text{G}\alpha$ should compete with $\text{G}\alpha\text{GTP}$ for interaction with p115 and block the observed GAP activity. As shown in Fig. 5, Panel A, GDP-AlF_4^- bound $\text{G}\alpha_{12}$ and $\text{G}\alpha_{13}$ effectively inhibited the GAP activity of p115 for $\text{G}\alpha_{12}$, while similar forms of $\text{G}\alpha_s$, $\text{G}\alpha_i$, and $\text{G}\alpha_q$ were without effect. Additionally, a titration of GDP-AlF_4^- bound forms of $\text{G}\alpha_{12}$ and $\text{G}\alpha_{13}$ demonstrated that the subunits are equipotent in inhibiting the GAP activity of $\text{G}\alpha_{13}$ (Fig. 5, Panel B). These competition assays suggest that the two G protein subunits have a similar affinity for p115 and supports the apparent differential efficacy of p115 towards the subunits as shown in Fig. 2.

Example 5. Binding of $\text{G}\alpha_{13}$ to p115 Rho GEF in vivo.

The following experiments demonstrated that $\text{G}\alpha_{13}$ and p115 Rho GEF interact in a GTP-dependent manner.

EXV-myc tagged (for COS cell transfections) and pAc-Glu tagged (for baculovirus expression) proteins with deletions of the RGS or DH domains were constructed as previously described in Hart et al., *J. Biol. Chem.*, **271**, 25452-25458 (1996). Full-length versions were constructed in the same vectors. A fusion of GST to the first 246 amino acids of p115 Rho GEF was constructed in pGEX4T-2 (Pharmacia). Transfections,

immunoprecipitations, and purifications were performed as previously described in Hart et al., *J. Biol. Chem.*, **271**, 25452-25458 (1996).

In COS cells transfected with myc-tagged p115 Rho GEF, $G\alpha_{13}$ can be specifically immunoprecipitated using the anti-myc antibody (Fig. 6, Panels A and B). This interaction is dependent on the presence of aluminum fluoride which is added to mimic the activated GTP-bound state of the $G\alpha_{13}$. Additionally, a truncated mutant of p115 Rho GEF which lacks the amino-terminal RGS domain is incapable of mediating co-immunoprecipitation, while full-length protein with a deletion in the DH domain does mediate co-immunoprecipitation. The differential binding of full-length and truncated Rho GEF proteins could also be detected using antibodies to $G\alpha_{13}$ to immunoprecipitate the complex (Fig. 6, Panel C). A very weak interaction with $G\alpha_{12}$ was detectable, while antibodies to $G\alpha_s$, $G\alpha_i$, $G\alpha_q$ and $G\alpha_z$ do not detect immunoreactive bands in the anti-myc immunoprecipitates, in spite of the fact that their respective antigens are detectable in the whole cell lysates. The co-immunoprecipitation of p115 Rho GEF and $G\alpha_{13}$ can be reproduced in a semi-purified system in which purified $G\alpha_{13}$ is added to immunoprecipitated p115 Rho GEF (Fig. 6, Panel D), suggesting a direct interaction. This direct interaction is consistent with the observation that p115 Rho GEF stimulates $G\alpha_{13}$ GTPase activity, but also indicates that p115 Rho may be an effector of $G\alpha_{13}$.

Binding could also be detected between the Rho GEF protein, KIAA380 and the α_{12} G protein subunit (Fig. 9, KIAA380 is referred to as FL147). In COS cells transfected with myc-tagged KIAA380, $G\alpha_{12}$ can be specifically immunoprecipitated using the anti-myc antibody (Fig. 9, Panels A and B, KIAA380 is referred to as FL147). This interaction is dependent on the presence of aluminum fluoride which is added to mimic the activated GTP-bound state of the $G\alpha_{13}$.

Example 6. Stimulation of p115 Rho GEF activity by $G\alpha_{13}$.

The ability of $G\alpha_{13}$ to affect the exchange activity of p115 Rho GEF was examined by incubating RhoA and p115 Rho GEF with or without $G\alpha_{13}$ to determine the effect on guanine nucleotide exchange.

RhoA (2.5 μ M) was loaded with [3 H]GDP by incubation at 30°C for 1 hour with 25 μ M GDP (10,000 cpm/pmol) in 50 mM NaHepes, pH 7.5, 50 mM NaCl, 4 mM EDTA, 1mM dithiolthreitol and 0.1% Triton X-100. After addition of MgCl₂ to 9 mM and octylglucoside to 1%, the Rho was incubated for an additional 5 minutes and separated from free GDP by rapid filtration through Sephadex-G50 that had been equilibrated with 50 mM NaHEPES, pH 7.5, 50 mM NaCl, 1 mM EDTA, 1 mM dithiolthreitol, 5 mM MgCl₂, and 1% octylglucoside. Dissociation of GDP from RhoA was measured at 30°C in 20 μ l of 50 mM NaHEPES, pH 7.5, 50 mM NaCl, 1 mM EDTA, 1 mM dithiolthreitol, 5 mM MgCl₂, 30 mM AlCl₃, 5 mM NaF, and 5 μ M GTP γ S. Unless specified, G protein alpha subunits were preincubated with AMF (30 μ M AlCl₃, 5 mM MgCl₂ and 5 mM NaF) prior to mixing with other proteins. Where indicated, alpha subunits were treated with 25 μ M GTP γ S or GDP β S rather than AMF and reactions were incubated without AMF but with 5 μ M of the respective nucleotide. Reactions were started with the addition of [3 H]-GDP-RhoA and bound GDP was determined by filtration (Northup et al., *J. Biol. Chem.*, **257**, 11416-11423 (1982)) prior to and after incubation.

The G α_s and G α_i alpha subunits were purified after expression in *Escherichia coli* (Lee et al., *Meth. Enzymol.*, **237**, 146-164 (1994)). The G α_q and G α_z alpha subunits were coexpressed in Sf9 cells with hexahistidine-tagged beta and gamma subunits and isolated as described (Kozasa and Gilman, *J. Biol. Chem.*, **270**, 1734-1741 (1995)). G α_{13} was prepared by a similar procedure to G α_{12} using baculovirus (Singer et al., *J. Biol. Chem.*, **269**, 19796-19802 (1994)) and octylglucoside during washing and elution of the α subunit after immobilization of the heterotrimer on Ni-NTA resin (Qiagen). The eluted G α_{13} was further purified by absorption to and elution from hydroxyapatite. About 500 μ g of purified G α_{13} can be obtained from 3 liters of cells. The expression of GST-RhoA in Sf9 cells, cleavage of the GST tag and isolation of the free RhoA were as described in Singer et al., *J. Biol. Chem.*, **271**, 4505-4510, (1996).

These studies demonstrated that the G α_{13} is capable of stimulating the activity of full-length p115 Rho GEF in a manner which depends on the concentrations of both p115 Rho GEF (Fig. 7, Panel A) and G α_{13} (Fig. 7, Panel B). The closely related alpha subunit

$G\alpha_{12}$ was ineffective in stimulating the activity of p115 Rho GEF in these experiments (Fig. 7, Panel A). Stimulation of Rho exchange was also monitored as a function of the activation state of $G\alpha_{13}$. The data graphed in Fig. 7, Panel C confirm that the stimulation of exchange activity is dependent on either aluminum fluoride (AMF) or $GTP\gamma S$, but is not stimulated by the deactivated nucleotide state mimicked by $GDP\beta S$. Additionally, a series of other alpha subunits including $G\alpha_q$, $G\alpha_z$, $G\alpha_s$, and $G\alpha_i$ also did not affect the activity of p115 Rho GEF (Fig. 7, Panel D). These results are consistent with the activated $G\alpha_{13}$ -dependent binding shown in Figure 6, and suggest that the productive binding of $G\alpha_{13}$ to p115 Rho GEF may be sufficient for activation.

Example 7. Effects of domains of p115 and $G\alpha_{12}$ on the p115 nucleotide exchange activity.

The theory that the RGS domain of p115 Rho GEF is normally autoinhibitory and that binding to $G\alpha_{13}$ relieves this inhibition was examined by comparing the effects of full-length Rho-GEF versus truncated Rho-GEF on Rho exchange activity.

Preparation of p115 proteins was as described in Example 1 above and as described in Hart et al., *J. Biol. Chem.*, **271**, 25452-25458 (1996). The assays shown in Figure 8, Panels B and C were performed as described in Example 5 above. AMF was the activating agent.

The results of these experiments showed that truncated p115 Rho GEF lacking the RGS domain demonstrates consistently elevated Rho exchange activity when compared with equal concentrations of the full-length protein (Fig. 8, Panel A). Additionally, addition of the isolated RGS domain (as a GST fusion protein) resulted in abrogation of $G\alpha_{13}$ -stimulated p115 Rho GEF activity (Fig. 8, Panel B). These data do not preclude additional $G\alpha_{13}$ -binding sites on p115 Rho GEF, although they do suggest a primary mode of action via the RGS domain.

The inability of the $G\alpha_{12}$ subunit to activate p115 Rho GEF was puzzling in light of the fact that p115 Rho GEF is capable of activating the GTPase of both $G\alpha_{12}$ and $G\alpha_{13}$. Therefore, an experiment was conducted in which $G\alpha_{12}$ was added to a $G\alpha_{13}$ -stimulated p115 Rho GEF assay (Fig. 8, Panel C). The results showed that $G\alpha_{12}$ was able to inhibit the

coupling of $G\alpha_{13}$ with p115 Rho GEF. This data is consistent with a model in which $G\alpha_{12}$ competes with $G\alpha_{13}$ for binding to the RGS domain of p115 Rho GEF. However, binding of $G\alpha_{12}$ to p115 Rho GEF is clearly not sufficient to stimulate Rho exchange activity. These results suggest that either the interaction of $G\alpha_{12}$ with the RGS domain of p115 Rho GEF is quite different from that of $G\alpha_{13}$ or that there may be an additional site of interaction between $G\alpha_{13}$ and p115 Rho GEF.

For other aspects of the nucleic acids, polypeptides, antibodies, etc., reference is made to standard textbooks of molecular biology, protein science, and immunology. See, e.g., Davis et al. (1986), *Basic Methods in Molecular Biology*, Elsevier Sciences Publishing, Inc., New York; Hames et al. (1985), *Nucleic Acid Hybridization*, IL Press, *Molecular Cloning*, Sambrook et al.; *Current Protocols in Molecular Biology*, Edited by F.M. Ausubel et al., John Wiley & Sons, Inc; *Current Protocols in Human Genetics*, Edited by Nicholas C. Dracopoli et al., John Wiley & Sons, Inc.; *Current Protocols in Protein Science*; Edited by John E. Coligan et al., John Wiley & Sons, Inc.; *Current Protocols in Immunology*; Edited by John E. Coligan et al., John Wiley & Sons, Inc. The entire disclosure of all patent applications, patents, and publications cited herein are hereby incorporated by reference.

From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

CLAIMS

What is claimed is:

1. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof,
5 consisting essentially of an RGS domain of a GEF protein.
2. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof,
comprising an RGS domain of a GEF protein, with the proviso that the polypeptide
does not comprise a DH domain or a PH domain.
10
3. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof,
wherein the polypeptide is selected from the group consisting of p115 Rho-GEF,
Lsc, KIAA380, and wherein the polypeptide is mutated in the RGS domain, and
wherein the polypeptide has a specific binding affinity for a G protein α subunit or a
15 GTPase activating activity for a G protein α subunit.
4. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof,
according to claim 1 or 2, wherein the GEF protein is a Rho GEF protein.
- 20 5. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof,
according to claim 4, wherein the Rho GEF protein is p115 Rho-GEF.
6. An isolated RGS-GEF polypeptide, or biologically active fragment thereof,
according to claim 4 wherein the Rho GEF protein is selected from the group
25 consisting of Lsc, KIAA380, and DrhoGEF2.
7. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof ,
according to claim 1 or 2, wherein the polypeptide has a specific binding affinity for
a G protein α subunit or a GTPase activating activity for G protein α subunits.

8. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof, according to claim 4, wherein the polypeptide has a specific binding affinity for a G protein α subunit or a GTPase activating activity for G protein α subunits.
- 5 9. An isolated RGS-GEF polypeptide, or a biologically active fragment thereof, according to claim 5, wherein the polypeptide has a specific binding affinity for a G protein α subunit or a GTPase activating activity for G protein α subunits.
- 10 10. An isolated RGS-GEF nucleic acid consisting essentially of a nucleotide sequence encoding a polypeptide comprising an RGS domain of a GEF protein.
11. An isolated RGS-GEF nucleic acid comprising a nucleotide sequence encoding a polypeptide comprising an RGS domain of a GEF protein, wherein the polypeptide does not include a DH domain or a PH domain.
- 15 12. An isolated RGS-GEF nucleic acid according to claim 10 or 11, wherein the GEF protein is a Rho GEF protein.
- 20 13. An isolated RGS-GEF nucleic acid according to claim 12, wherein the Rho GEF protein is p115 Rho GEF.
14. An isolated RGS-GEF nucleic acid according to claim 12 wherein the Rho GEF protein is selected from the group consisting of Lsc, KIAA380, and DrhoGEF2.
- 25 15. An isolated RGS-GEF nucleic acid according to claim 10 or 11, wherein the polypeptide has a specific binding affinity for a G protein α subunit or a GTPase activating activity for a G protein α subunit.
- 30 16. An isolated RGS-GEF nucleic acid according to claim 12, wherein the polypeptide has a specific binding affinity for a G protein α subunit or a GTPase activating activity for a G protein α subunit.

17. An isolated RGS-GEF nucleic acid according to claim 13, wherein the polypeptide has a specific binding affinity for a G protein α subunit or a GTPase activating activity for a G protein α subunit.
- 5 18. A method of modulating an activity of a G protein α subunit comprising, administering to a mammal an effective amount of a polypeptide according to claim 1 or 4.
- 10 19. A method of identifying or assaying a molecule that inhibits or enhances binding of a monomeric G protein guanine nucleotide exchange factor to a G protein α subunit comprising incubating the G protein α subunit, or fragments thereof, with the monomeric G protein nucleotide exchange factor, or fragments thereof, in the presence and absence of a test molecule and determining whether the presence of the
- 15 test molecule inhibits or enhances binding between the monomeric G-protein guanine nucleotide exchange factor and the G protein α subunit.
20. A method of identifying or assaying a molecule that inhibits or enhances a stimulatory effect of a GEF on a G α subunit GTPase activity comprising incubating
- 20 a G α subunit, or fragments thereof, with a GEF protein, or fragments thereof, in the presence and absence of a test molecule and determining whether the presence of the test molecule inhibits or enhances the stimulatory effect of the GEF protein on G α subunit GTPase activity.
- 25 21. A method of identifying or assaying a molecule that specifically inhibits the stimulatory effect of an activated G α subunit on GEF mediated nucleotide exchange of a monomeric G protein, comprising conducting a first assay by incubating an activated G α subunit, or fragments thereof, with a GEF protein, or fragments thereof, and a monomeric G protein, or fragments thereof, in the presence and
- 30 absence of a test inhibitor, conducting a second assay by incubating a GEF protein,

or fragments thereof, and a monomeric G protein, or fragments thereof, in the presence and absence of the test inhibitor, and determining whether any inhibitory effect of the test inhibitor in the first assay is greater than any inhibitory effect of the test inhibitor in the second assay.

5

22. A method of identifying or assaying a molecule that specifically enhances the stimulatory effect of an activated G α subunit on GEF mediated nucleotide exchange of a monomeric G protein, comprising conducting a first assay by incubating an activated G α alpha subunit, or fragments thereof, with a GEF protein, and fragments thereof, and a monomeric G protein, or fragments thereof, in the presence and absence of a test enhancer, conducting a second assay by incubating a GEF protein, or fragments thereof, and a monomeric G protein, or fragments thereof, in the presence and absence of the test enhancer, and determining whether any enhancing effect of the test enhancer in the first assay is greater than any enhancing effect of the test enhancer in the second assay.

15

23. A method of identifying or assaying a molecule that mimics the stimulatory effect of an activated G α subunit on GEF mediated nucleotide exchange of a monomeric G protein comprising identifying a test compound that exhibits a binding affinity for the RGS domain of GEF proteins, or fragments thereof, incubating a GEF protein, or fragments thereof, and monomeric G protein, or fragments thereof, in the presence or absence of the test compound, determining whether the test compound exhibits a stimulatory effect on GEF mediated nucleotide exchange of a monomeric G protein.

20

24. A method of identifying or assaying a molecule that mimics the stimulatory effect of an RGS domain of a GEF protein on GTPase activity of a G α subunit comprising identifying a test compound that exhibits a binding affinity for a G α subunit and incubating a GTP loaded G α subunit in the presence or absence of the test compound to determine whether the test compound has a stimulatory effect on G α subunit GTPase activity.

30

25. A method according to claim 19, 20, 21, 22, 23, or 24 wherein the GEF protein is selected from the group consisting of p115 Rho GEF, Lsc, KIAA380, and DrhoGEF2.
- 5
26. A method of expressing in transformed host cells, a polypeptide coded for by a nucleic acid, comprising culturing transformed host cells containing a nucleic acid according to claim 11.
- 10
27. A transformed cell containing a nucleic acid according to claim 11.
28. A vector comprising a nucleic acid according to claim 11.

```

RGS4 2nd St.      *          **          *
RGS4 (137)   T S R N M L E - - P T I T C F D E A Q K K I F N L M E K D S Y R R F L K S R F Y L D (176)
RGS2 (158)   I A Q N I Q E - - A T S G C F T T A Q K R V Y S L M E N N S Y P R F L K S E F Y Q D (197)
GAP (165)    I N K M Q E - - P S A H T E D D A Q L Q I Y T L M H R D S Y P R F L K S S P T Y R A (204)
RGS12 (792)  D D I L N A - - P H P D M F K E Q Q L Q I F N L M K F D S Y T R F L K S Q L Y Q E (830)
RGS14 (144)  E E V L A Q - - P R P D M F R A Q Q L Q I F N L M K F D S Y A R F V K S P L Y Q E (182)
p115 (130)   I S E D V Q - R R F V Q E V V Q S Q Q V A V G R Q L E D F R S K R L M G M T P W E Q (170)
Lac (128)    I S E D V Q - R R F I Q E V V Q S Q Q A A V S R Q L E D F R S K R L M G M T P W E Q (168)
XIAA380 (394) - S E D A - R G V L C E A Q E A A M P E I S E Q L I H D Y R T K R T I G L G S L Y G (432)
DrhoGEF2 (1012) Y D K V E I L R T V F L R S R K R A K D L I S E Q L R E F O O K R T A G L G T I Y G (1053)

```




FIGURE 1

2/34




FIGURE 2

3/34




FIGURE 3

4/34




FIGURE 4

5/34




FIGURE 5

6/34




FIGURE 6

7/34




FIGURE 7

8/34




FIGURE 8

9/34




FIGURE 9

SUBSTITUTE SHEET (RULE 26)

10/34

10	20	30	40
medfargaaspgpsrpglvpvsii gaededfeneletnse 40			
eqnsqfqsleqvkrpahlmallqhvalqfepgpilccih 80			
admlgslgpkeakkafldfyhsflectavlrpvppnvaf 120			
eldrtradlisedvqrrfvqevvqsqqvavgrqledfrsk 160			
rlmgmtpweqelaqlaawvgrdrasyeareerhvaerllmh 200			
210	220	230	240
leemqhtistdeeksaavvnaiglymrhlgvrtksgdkks 240			
grnffrkkvmgnrrsddppktkkglssil daarnwngepq 280			
vpdfrhlkaevdaekpgatdrkggvgmpsrdrnigapqad 320			
tpgvsllhplsl dspdrepgadaplel gdsspaggpmslesl 360			
appestdegaetespepgdegepggrsglelepeppgwre 400			
410	420	430	440
lvppdtl hslpksqvkrqevise llvteaahvrmlrvlhd 440			
lffqpmacel ffp leelqni fpsl delievhs lfldrlmk 480			
rrqesgyll eeigdvll arfdgaegswfqkissrfcsrqs 520			
faleqlkakqrkdprfcavqeaesrprcrrlqlkdmipt 560			
emqrltkyp lllqsigqnteepterekvelaaeccreilh 600			
610	620	630	640
hvnqavrdmedllrlkdyqrrldlshlrqssdpmlsefkn 640			
lditkkklvhegpltwrvtkdkavevhvllldlllllqr 680			
qderlllkshsrtltptpdgktmlrpvlrltsamtrevat 720			
dhkafyvllftwdqeaqiyelvaqtvserknwcalitetag 760			
slkvpapasrpkprrpsstrep lssengnggretspa 800			
810	820	830	840
darteril sdl l pfcprpgpegqlaata l rkvislkqllfp 840			
aeedngagpprdgdgvpggplspartqei qenllsleet 880			
mkqleeleeeefcrlrp llsq l ggnsvpapgct 912			

FIGURE 10

10 20 30 40

gggcgccccgcgggtcacttccgcgcggacaccagccttg 40
cagagcccagggagatggaagacttgcgccgaggggcggc 80
ctccccaggccccctcccggcctggcctggttcccgtcagc 120
atcatcggggctgaggatgaggattttgagaacgagctgg 160
agacaaactcagaagogcaaaacagccagttccagagcct 200

210 220 230 240

ggagcaggtgaagcggcgcccagcccacctcatggccctc 240
ctgcagcacgtggccctgcagtttgagccaggaccctgc 280
tttgctgtctgcattgccgacatgctgggctcactgggccc 320
caaggaggccaagaaggccttccctggacttctaccacagc 360
ttcctggagaagacagcgggttctccgggtgcccgtccctc 400

410 420 430 440

ccaacgtcgctttgaacttgaccgcactagggctgacct 440
catctccgaggatgtccagcggcggttcgtgcaggaggtg 480
gtgcaaagccagcaggtagccgtgggcgggcagctggagg 520
acttccgttccaagcggctcatgggcatgacgccctggga 560
gcaggagctggcccagctggaggccttgggttgggcgggac 600

610 620 630 640

cgagccagctacgaggccccgggagcggcacgtgycgggagc 640
ggctgctcatgcacctggaggagatgcaacataccatctc 680
taccgacgaagaaaagagtgcctgccgtgggtcaatggccatt 720
gggctgtacatgcgccaccttgggggtgcggaccaagagtgc 760
gagacaagaagtcggggagggaacttcttccggaaaaaagggt 800

810 820 830 840

gatggggaaccggcggtcggacgacctcccaagaccaag 840
aaggggctgagcagcatcctggatgccgcccgcctggaacc 880
ggggagagccccaggttccagattttcgacacctcaaagc 920
agagggtgatgccgagaagccaggtgctacagaccggaag 960
ggaggcgtggggatgccctctcgggaccggaatatcgggg 1000

1010 1020 1030 1040

ctcctgggcaggacacccctggagttctctctgcacctct 1040
gtccctggacagcccagaccgggaaccaggtgctgacgcc 1080
cccctggagctgggggactcatccccgcaggggcccaatga 1120
gcctggagtccttggcgccccccagagagtaccgacgaggg 1160
ggccgaaccgagagccccgagcctggagatgaggggggag 1200

1210 1220 1230 1240

ccggggcggtcgggactggagcttgaaccagaagagcctc 1240
ccggctggcggggaactcgtccccccagacacctgcacag 1280
cctgcccgaagagccaggtgaagcggcaggaggtcatcagc 1320
gagctgctggtgacagaggcgcccacgtgcgcattgctgc 1360
gggtgctgcacgaccttcttcttccagcccatggcagaatg 1400

SUBSTITUTE SHEET (RULE 26)

12/34

1410	1420	1430	1440
.....
cctgtttcttcccccttggaggagctgcagaacatcttcccc	1440		
agcctggacgagctcatcgaggtgcattccctgttcctcg	1480		
atcgctgatgaagcggaggcaggagagtggctacctcat	1520		
cgaggagatcgagagacgtgctgctggcccggtttgatgg	1560		
gctgagggctcctggttccagaaaatctcctcccgccttct	1600		
1610	1620	1630	1640
.....
gcagccgccaagtcatTTgccttagagcagctcaaagccaa	1640		
gcaacgcaaggaccctcggttctgtgccttcgtgcaggaa	1680		
gctgagagccgcccgcggtgccgcccgtgcagctgaagg	1720		
acatgatccccacggagatgcagcggctgaccaagtaccc	1760		
cctgctcctgcagagcatcgggcagaaacacagaagagccc	1800		
1810	1820	1830	1840
.....
acagaacgggagaaagtggagctggcagccgagtgctgcc	1840		
gggaaattctacaccacgtcaaccaagccgtgcgtgacat	1880		
ggaggacctgctgaggctcaaggactatcagcggcgccctg	1920		
gacttgctccacaccttcggcagagcagcgaccctatgctga	1960		
gcgagttcaagaacctggacatcaccaagaagaatttgg	2000		
2010	2020	2030	2040
.....
ccacgagggcccaactgacgtggcgggtgactaaggacaag	2040		
gcagtggagggtgcatgtgctgctgctggacgacctgctgc	2080		
tgctgctccagcgccaggacgagcggctgctgctcaagtc	2120		
ccatagccggacactgacgcccacgcccgatggcaagacc	2160		
atgctgcggcccgtgctgcggctcacctccgccatgaccc	2200		
2210	2220	2230	2240
.....
gcgaggtggccaccgatcacaaagccttctacgtcctttt	2240		
tacctgggaccaggaggccagatatagagctgggtggca	2280		
cagactgtgtcggagcggaaaaactgggtgtgctctcata	2320		
ctgagactgccggatccctgaaagtccctgcccctgcctc	2360		
tcgccctaagccccggcccagggcggagcagcaccgagaa	2400		
2410	2420	2430	2440
.....
cccctcctcagcagctctgagaacgggaatggtggccgag	2440		
agacgtctccagctgatgcccggaaccgagagaatcctcag	2480		
tgacctcctgccccttctgcagaccaggccccgagggccag	2520		
ctcgctgccacggccccttcggaaagtgtgtccctgaagc	2560		
agcttctgtttccggcggaggaagacaatggggcggggcc	2600		
2610	2620	2630	2640
.....
tcctcgagatggggatgggggtcccagggggcggggcccctg	2640		
agcccagcacggaccaggaatccaggogaacctgctca	2680		
gcttggaggagaccatgaagcagctggaggagtggaggga	2720		
ggaaattttgccgcctgagocccctcctgtctcagcttggg	2760		
gggaactctgtccccccagcctggctgcacttgagggttccc	2800		

FIGURE 11 cont.

13/34

2810	2820	2830	2840
<hr/>			
gcccaggaaggccttttgcaagaaggagaggaatggggga 2840			
gaggacgtgagggaccacccccacccacacagctgccgca 2880			
gcatctcacacccccgagggcctgaggagagggagctgtgg 2920			
gccacgcctgggagggggcccagctgggggttactgcccccg 2960			
catgagcctcggccatctctccctcctgccctctgcttgg 3000			
3010	3020	3030	3040
<hr/>			
gggactcagggctccattctggagggcaccacggtgaccc 3040			
gggccatctcagtattgcctgtgggggccacccctccacc 3080			
cccccccccaagtgccttcgctctgtttttataccctgaa 3120			
ttggagggtttatttttttaatatatattat 3150			

FIGURE 11 cont.

SUBSTITUTE SHEET (RULE 26)

14/34

10	20	30	40
msvrlpqsidrlsslsslgdsaperkspshhrapsdaset 40			
tglvqrcviiqkdqhgfgftvsgdrivlvqsvrpggaamk 80			
agvkegdriikvngtmvtntsshlevvklksgayvaltll 120			
gsspsmgisglqqdpspagapritsvipspppppplppp 160			
qritgpkplqdpevqkhatqilrnmirqeekelqdiiply 200			
210	220	230	240
gdtqrpsegrlslsdsqegdsqldsgterfpslseslmnr 240			
nsvlsdpqlspsrtspvimarvaqhrrragsdaavpstgd 280			
qgvdaqspkplliigpeedydpgyfnnesdiiifqdleklksr 320			
pahlgvflryifsqadpspllfylcaevyqaaspkdsrsi 360			
gkdiwnifleknapiirvkipemlqaeidsrlrnsedargv 400			
410	420	430	440
lceaqaampeiaeqeihdyrtkrtlglslygendlildid 440			
gdplrerqvaekqlaalgdilskyedrsapmdfalntym 480			
shagirlrearpstntaekaqsapdkdkwlpffpktkkssn 520			
skkekdaledkkrnpilkyigkpksssqstfhiplspvev 560			
kpgnvrniiahfennqaydaepqgtarlstgsfpedilles 600			
610	620	630	640
dssrseirlgrsesikgreemkrsrkaenvprsrsvdmd 640			
aaeatrlhqsassstsslstrslenptppftpkmgrrsi 680			
espslgfctdtllphlledldgqlsdlepepdaqnwqhtv 720			
gkdvvagltqreidrquevinelfvteashlrtlrvldlif 760			
yqrmkkenlmpreelarlfpnlpelieihnsweamkkir 800			
810	820	830	840
eegpiikeisdmlarfdgpareelaqvaafcsyqsiat 840			
eliktqarkesrfqlfmqeaeshpqcrrlqlrdliisema 880			
rltkyplllesiiikhteggtseheklcrardacreilkv 920			
neavkatenrhrlegyaqrldatalerasnplaaefksid 960			
lttrkmihegpltwriskdktldihvllledllvllqkd 1000			
1010	1020	1030	1040
eklllkchsktavgsdskatfspvklnavlirsvatdk 1040			
raffiictsklgppqiyei valtssdkntwmelleavrn 1080			
atrhpgaampvhhppppgprepaqagptpsrvelddsdvf 1120			
hgepepeelpggtgsqarvaghqvlledpeqegsaeeee 1160			
lgvlpcpstslngenrgirtrnpihlafpgplfmeqlads 1200			
1210	1220	1230	1240
aledvenlrhlilwslilpghtmetqaaqepeddltppsv 1240			
isvtshpwpdpgspgqappgggedntqlaglegerpeqedm 1280			
glcslahlpprrtrnsgiwespeidrnl aedassteaaggy 1320			
kvvrkaevagskvvpalpesgasepgppeveggtkatgnc 1360			
fyvsmppsgppdssthdseapmsppapdsipagqtepapqi 1400			

FIGURE 12

15/34

1410 1420 1430 1440

qggnddprprsrpspslalrdvgmifhtieqltklnrik 1440
dmeiahrellkslggessggttpvgsfhteaarwdgsls 1480
ppakeplasdrrnsheigpcpedgsdapledstadaasp 1520
gp 1522

FIGURE 12 cont.

SUBSTITUTE SHEET (RULE 26)

16/34

10	20	30	40
aattggctcatttaagaatttcaaaacattttaatgtaaaa 40			
gcttttttttttttaaggaagtccataaattttggttccc 80			
agggttgcaactggacttggaaggagtgctgttggtacat 120			
actattgtatggttttattttatttttactgtacaaatc 160			
agccgaaagaatttttccaagtgccattttcggattttatta 200			
210	220	230	240
atccttttttttttccctttccctcaaagatatittgctgttg 240			
tcatattaagcattggagactagaaaattactttcccccct 280			
ttgagctagaggggtctcttgccaacagaaggacagctgag 320			
aaagctggatttaaaaggatggttttatctgtactttgcag 360			
ttaacagtgotattttgaaggcacatttttctgtgattca 400			
410	420	430	440
tttttttttgccatagtgttaaccttgaagagatttcgtg 440			
gctgggttttttggtttctgagaaggctcgtagtttttctc 480			
ttttccttttttttttcttttttcttttcttttctttt 520			
ttttaaagcgggggaggggaaggaggggctgagaaaggaaa 560			
tcatgttcaactggtagaagttagagtggagcatcagttacc 600			
610	620	630	640
agggctcctgogagctggaggcgaaaggatttctatcttcaa 640			
gttgggaggccctcctctcaccttgctcaaaaattgcaag 680			
cgattcaatcctgatcaagacaccaaaagctacaggattct 720			
ggaaccgtggagacaccgagaaaccatgagtgttaagggtta 760			
ccccagagtatagacagggttaagtagcctgtcttctcttg 800			
810	820	830	840
gagattctgcaccagagcgcaagtcccccttcccaccatcg 840			
ccagccttcggatgcctctgagacaacagggtctcgttcaa 880			
cgctgtgtcattatccaaaaggaccagcatggcttcggct 920			
tcacagtcagtggggatcgcatgttcttggtgcagtcgtgt 960			
gcggcctggagggtgcagccatgaaggccggtgtgaaagag 1000			
1010	1020	1030	1040
ggcgaccggatcatcaaagtcaacggcaccatggtgacca 1040			
atagctcacacctggaagtggtaaaagctgatcaaatctgg 1080			
cgcctatgtcgcactcaccctcctgggctcttcaccttca 1120			
tccatgggcattctctgggctccagcaggaccatccccag 1160			
caggagctccccgaatcacgtcagtgatccccctcaccacc 1200			
1210	1220	1230	1240
acctcctccacctctaccacctccacaacgcatacacagga 1240			
cccaaacctctgcaggatcccgaagttcaaaaacatgccca 1280			
cccagatcctcaggaatatgctgaggcaggaagaaaaaga 1320			
attacaggacatacttccactatatggtgacaccagccag 1360			
agaccatcagaaggccggctctctcttggttcccaggagg 1400			

FIGURE 13.

1410 1420 1430 1440

gggacagtggccttggaactctgggacagaacgctttccttc 1440
cctcagtgagtcattgatgaatcggaactcggtaactgtca 1480
gaccctgggctagacagtcctcgaaacctcccctgtgatca 1520
tggccaggggtggcccagcaccacaggcgagggtcggga 1560
tgcagcagtcacctcaaccgggtgaccaggggtgtagatcaa 1600

1610 1620 1630 1640

agcccaaagcctttaattattggcccagaggaagactatg 1640
acccggttatttcaacaacgagagcgacatcatattcca 1680
ggatctgggagaaactgaagtctcggccagctcacctgggg 1720
gtttttctacgttacatcttctctcaggcggaacccagtc 1760
cactgcttttttacctgtgtgcagaagtttatcagcaggc 1800

1810 1820 1830 1840

aagccccaaggattcccgaagcttggggaaagacatctgg 1840
aatattttcctggagaaaaatgcgcctctgagagtgaaga 1880
tcctgagatgctacaggctgaaattgactcgcgcctgcg 1920
gaacagcgaagatgcccgtaggtgttctctgtgaagctcaa 1960
gaggcagccatgcctgagatccaagagcagatccacgact 2000

2010 2020 2030 2040

acagaacgaagcgcacactggggctgggcagcctgtatgg 2040
tgaaaatgacctgctggacctggatggggacctctccga 2080
gagcgccaagtggctgagaagcagctggctgcccttggag 2120
atatittgtccaagtatgagggaagacaggagcgcacctat 2160
ggacttcgccctcaatacctacatgagccatgctgggac 2200

2210 2220 2230 2240

cgtcttcgagaggcacgaccttccaacacagctgaaaagg 2240
cccagtctgctcctgacaaggacaagtggctaccgttctt 2280
ccctaagaccaagaagagcagcaattccaagaaaagaaag 2320
gatgccttggaggacaagaagcgaaacctatcctcaaat 2360
acattgggaagcccaaaagctcttctcaaaagcacatttca 2400

2410 2420 2430 2440

tattcccttgtccccctgtggaagtcaaaccaggcaatgtg 2440
aggaacatcattcagcactttgagaacaaccagcagtatg 2480
atgccccagaacctgggacacaacgactctcgaccgggaag 2520
ctttcctgaggacctgctggagagtgacagttcacgctca 2560
gagattcgcttggggcgctctgaaagcctcaaggggccggg 2600

2610 2620 2630 2640

aagagatgaaacggtctcgaaggcagagaacgtgccccg 2640
ctctcgcagtgatgttgacatggatgctgctgcggaggct 2680
actcgctgcaccagtcagcctcgctcctctacctccagcc 2720
tctccaccaggctctcttgagaacccaacccctccattcac 2760
tcccaaaatggggccgcaggagcattgagtcccccagtttg 2800

SUBSTITUTE SHEET (RULE 26)

18/34

2810	2820	2830	2840
.....
gggttctgcacagataccctccttccccacctcctagagg	2840		
atgatctgggccagctgtctgacctggagccagagccaga	2880		
tccccaaaattggcagcatacagtgggcaaggatgtggtg	2920		
gctgggctaaccagcgaggagattgaccggcaagaggta	2960		
tcaatgagctgtttgtgactgaagcttcccacctgcgcac	3000		
.....
3010	3020	3030	3040
.....
actccgggtcctggacctgatcttctaccagcgaatgaag	3040		
aaggagaacctgatgccccgggaggagctggccccggctct	3080		
tccgaacctgcctgaactcatagagattcacaattcctg	3120		
gtgtgaagccatgaagaagctccgggaggaaggcccatc	3160		
atcaaagagatcagtgacctcatgctggccccggtttgatg	3200		
.....
3210	3220	3230	3240
.....
gccctgcccggagaggaaactccagcaagtggctgcacagtt	3240		
ctgttcctatcagtcgaatagccctagagctaataagacc	3280		
aagcaacgcaaggagagtcgattccagctcttcatgcagg	3320		
aggctgagagccaccctcagtgctcgcggtgtagctgag	3360		
agacctcatcatctctgagatgcagcggctcaccaagtac	3400		
.....
3410	3420	3430	3440
.....
ccgctgctgctggagagcatcatcaagcacacagagggtg	3440		
gcacctctgagcatgagaagctgtgccggggccgggacca	3480		
gtgccgggagattctcaagtatgtgaatgaagcggtaaaa	3520		
caaacagagaaccgcccaccgtttagagggctaccagaaac	3560		
gcctggatgccaccgcccctggagagggccagcaacccccct	3600		
.....
3610	3620	3630	3640
.....
ggcagcagagttcaagagcctggatcttacaaccagaaaa	3640		
atgatccatgagggacccctgacctggaggatcagcaagg	3680		
ataagaccttggacctccacgtgctgctgctggaggacct	3720		
cctagtgtgctacagaaacaggatgagaagctattgctg	3760		
aagtgccacagcaagactgctgtgggtcctcagacagca	3800		
.....
3810	3820	3830	3840
.....
agcagaccttcagccccgtgctcaagctcaatgctgtgct	3840		
catecgctctgtggccacagataaacgggccttcttcate	3880		
atctgcacctccaagctgggcccaccccagatctatgagc	3920		
tggttgcattgacgtcatcagacaagaacacatggatgga	3960		
gctcttagaagaggccgtgcggaatgccaccaggcacccc	4000		
.....
4010	4020	4030	4040
.....
ggagctgccccaatgcccgctccatcctccacccccaggtc	4040		
cccgggagccagcccagcaggggccccacaccagcagggt	4080		
agaactggatgactcagacgtgttccatggtgaacctgaa	4120		
cctgaggagctgcctggaggcactgggtcccagcagaggg	4160		
tccaagggaagcaccagggtcctgctagaggacctgagca	4200		

FIGURE 13 cont.

19/34

4210	4220	4230	4240
gaggggagtgagaggaagagggaactgggtgtcctgct			4240
tgcccttccacatccctggatggagagaacaggggcatca			4280
ggacaaggaaccccatccacttggccttcccaggccctct			4320
gttcatggaagggtcgctgactccgctctggaagatgtg			4360
gagaacctgcgacatctgatccctgtggagcctgctgccag			4400
4410	4420	4430	4440
gtcacaccatggaaactcagggtgcccaggagcccaggga			4440
cgacctgacacccacaccttctgtcatcagcgtcacctct			4480
caccttgggacccagggtccccagggaagcaccacctg			4520
ggggtgaaggggacaacacccagcttgcagggtggagg			4560
ggaacggccagagcaggaagacatgggtctctgttctctg			4600
4610	4620	4630	4640
gaacacctaccccccaaggaccagaaattctgggatatggg			4640
agttctccagaactggacaggaatctggctgaagatgcttc			4680
aagcacagaggcagcaggaggttacaagttgtgagaaaa			4720
gctgaggtggcaggcagcaaggttgccctgcactaccag			4760
agagtggccagtcagagcctgggccacctgaagtgggaagg			4800
4810	4820	4830	4840
cggaacaaagggtacggggauactgcttttatgtcagcatg			4840
ccatcaggacccccggactcaagcaccgaccactcagagg			4880
cacctatgagccccctcagcctgacagcctccctgcagg			4920
gcagacagagcctcagcctcagctgcaggagggaacgat			4960
gatccaagacgccccagccgctctcctccaagcctggccc			5000
5010	5020	5030	5040
tcaggggacgtgggcatgatcttccatoccatgtgagcagct			5040
cactctcaagctcaacagggtcaaggatatggagctggcc			5080
cacagagagctgtcgaagtccttggggggagagtcacctg			5120
gtggcaccacgcctgtgggcagtttccacacagaagcagc			5160
tagatggacagatggctccctctcacctcccgcctaaggag			5200
5210	5220	5230	5240
cccctagcttctgactccaggaacagccatgaactggggc			5240
cctgccctgaggatggctctgacgccccctgggaagacag			5280
cacagcagacgcagccgcgtcaccaggaccataaccgtac			5320
aaaccaccaaatcctctgcgtccccactcctccttcagg			5360
actggcctgagacccggggcacagggtagggggggatcccaa			5400
5410	5420	5430	5440
cactcctccctgtggaggaggcagttagggaactaggat			5440
ccagccaaggcccggggggagacccgcagttgtcttggtc			5480
tgctcaagtcggagtcagggtttcagtgcttttccctccc			5520
ttagcccaaccttccaaggcctcatgtctcctaagcatgc			5560
tgactgcattccgaaggccccccactcaccatgggtctgcc			5600

FIGURE 13 cont.

SUBSTITUTE SHEET (RULE 26)

20/34

5610	5620	5630	5640
<hr/>			
tcacccacatatgtgtgtacacgcgcacgcctgtatgtg	5640		
cgctgccctcagacatgcaagtgaaggaggaggcttctg	5680		
tgtaaatgcactttcttctccctctttctccataagac	5720		
cccaggcagagggtgggtgcctccctccctctttgtcac	5760		
tttggtttcctataaatatgtatgtatcgt	5790		

FIGURE 13 cont.

21/34

10	20	30	40
mgevaggaapgprrsglvsiigaaeddfeneleansedq 40			
nsqfqsleqvkrpahlmallqhvalqfepgpilccihad 80			
mlsslgpkeakkafldfyhsflectavlrpvppsvafel 120			
drtrpdlisedvqrrfiqevvqsqaavsrqledfrskrl 160			
mgmtpweqelsilepwigkdrngyearerhvaerllshle 200			
210	220	230	240
etahtistdeeksaavvtailsymrhlgvrtksgdkksg 240			
nffrkkvmgnrrsdeppkttkglssildparwnrgesap 280			
dcrhikveadekpgpadrkgglgmssrdrtvgtpgqanpg 320			
vsihplstdsvdsrepgvdtppqepgdtppagptsleplap 360			
pestedngetespepgddgepggrsglelepeeppgwrelv 400			
410	420	430	440
ppdtllslpksqvkraqevisellvteaahvrmlrvlhdif 440			
yqpmadggffpldelqnfpsldelievhsifldrimkrr 480			
qesgylieeigdvllarfdaegswfqkissrfcsrqsfa 520			
leqlkakqrkeprfcavqeaesrprcrrlqlkdmiptem 560			
qrltkyplllqsigqnteestergkvelaaeccreilhhv 600			
610	620	630	640
nqavrdmedllrikdyqrrldlthlrqssdpmlsefknlid 640			
itkkklvhhegpltwrvtkdkaievhlldlllllqraqd 680			
erlllkshsrtltptpdgktmirpvlrltsamtrevatdh 720			
kafyviftwdqeaqiyeivaqtsserknwnlittetagsl 760			
kvpapasrlkprpsspsirepllssengtggamapada 800			
810	820	830	840
rterllndllpfcrrpgpegqlaatalqkvlsikqillste 840			
edsgagpprdgdgvpgrappgvhtqeieenllslevair 880			
qleeleefcrlrpillsqlggtlspnlaapersaqtgls 919			

FIGURE 14

22/34

10 20 30 40

cgggacaccgggcccgggatccccgagcccgacctcgggc 40
gcctcgccggtcacctccccgcgcggacaccagccccggcg 80
gagtcctggagatgggagaagtgcgcggagggggcggcccc 120
agggcctccccggtcttgccctggtgtccatcatcatcggg 160
gcggagggatgaggattttgagaacgagctggaggcgaaact 200

210 220 230 240

cagaagatcaaaacagccagttccagagcctagagcaagt 240
gaagcgccgccccctgcccacctcatggccctcctgcagcat 280
gtggccctgcagttcgagccaggaccactgctctgctgcc 320
tgcatgcagacatgctgagctctcttgggcccccaaggaagc 360
caagaaggccttccttgacttctatcacagtttccctggag 400

410 420 430 440

aagactgcggttctacgggtgcccgtccctcccagtgctcg 440
cttttgaacttgatcgtactcgacctgatctgatctctga 480
ggatgtccagaggcggttcatacaagagggtggtgcagagc 520
cagcaggcagccgtgagccgtcagctagaggacttccgct 560
ccaagcggctcatgggcatgacgccctgggagcagggaact 600

610 620 630 640

gagcctgctggagccctggaattgggaaagaccgagggaac 640
tatgaggccccgggagcggcatggtgcggagcggctgctgt 680
cccacctggaggagagcccagcataccatctctacagatga 720
agagaaaagtgctgctgtgtggtcactgccatcagcctgtat 760
atgcgccaccttggaagtccggaccaagagtggggacaaga 800

810 820 830 840

agtccgggaaggaaacttcttccggaaaaagggtgatggggaa 840
tcggagggtcagacgaacccccaaagacaaagaaagggctg 880
agcagtatcctagatcctgcacgttggaaaccggggagagc 920
catccgctccagattgtcgacatctaaagggtcgaggctga 960
tgagaagccaggccctgcagaccggaaggggaggcctgggt 1000

1010 1020 1030 1040

atgtctttctcgggacaggactggttgggactcctggacagg 1040
acaacccaggagtgcttccctgcacctctgtctacagacag 1080
cgtcgactcccgggaaccaggcggtggataccccgcaggag 1120
ccagggggatacacccccacaggggccctaccagcctggagc 1160
ccctggcgccccccagagagcacagaggacaatggcgagac 1200

1210 1220 1230 1240

tgagagccctgagcccggagatgatggggagccaggacgg 1240
tcgggcctggaaactggaaccagaagaacctcctgggtgga 1280
gggaactcgtgccccacagacacctgctcagttctgcccac 1320
gagccaaagtgaagcggcaagagggtcatcagcgagctgctc 1360
gtgocctgaggcagctcacgtgcgcatgctacgggtactgc 1400

FIGURE 15

23/34

1410	1420	1430	1440
atgacctcttctaccagcccatggcggtggaggcttctt	1440		
ccctctggacgagctgcagaacatcttcccagacctggat	1480		
gagctcatcgagggtgacctccctgttcctcgatcgcttga	1520		
tgaagcggagacaagagagtggtacctcattgaggagat	1560		
cggcgatgtgctactggccccggttcgatggtgctgagggc	1600		
1610	1620	1630	1640
tcattggttccagaagatctcctcccgttctgcagccgcc	1640		
agtcgttcgctctagagcagctcaaagccaagcagcgcaa	1680		
ggagcctcggttctgtgcctttgtgcaggaagctgagagc	1720		
cggccgagatgccggcgccctacagttaaaggacatgatcc	1760		
ccactgagatgcagcgactgaccaagtacccactgctgct	1800		
1810	1820	1830	1840
acagagcctcgggcagaaacacagaggagtctacagaacga	1840		
gggaagtgagcttgagctgagtgctgccgggagattc	1880		
tgcaccatgtcaatcaagccgtccgtgacatggaggacct	1920		
gctgcggctcaaggattaccagcggcgccctggacttgact	1960		
cacctacggcagagcagtgaccctatgctgagcgagttca	2000		
2010	2020	2030	2040
agaacctggacatcactaagaagaagttggtcatgaagg	2040		
ccccctcacgtggcgagtgaccaaagacaaaagctatagaa	2080		
gtgcacgtgctcttgctggacgacctgctgctgctgcc	2120		
agcgccaggacgagaggctgctgctcaagtcccacagccg	2160		
gacgctgacacctacccccgatggcaagaccatgctgcgg	2200		
2210	2220	2230	2240
ccggtgctccggctcacctctgccatgacccgagaggtagg	2240		
ccactgatcacaaagctttctacgtcatttttacctggga	2280		
ccaggaggcccgatatatgagctggtggcacagacatct	2320		
tcggaaacgcaaaaactggtgtaacctcatcactgagactg	2360		
ctggatccctgaagggtccctgccccctgcctcccgcctcaa	2400		
2410	2420	2430	2440
accccgccccagcccaagcagcatccgagaacccctgctc	2440		
agcagctctgagaatggcactggaggcgagagatggctc	2480		
cagctgatgccaggacagagcggctcctcaatgacctcct	2520		
gcccttctgcagaccaggccccagaggccagcttgctgcc	2560		
acagcccttcagaaagtactgtccctgaagcagatcctgc	2600		
2610	2620	2630	2640
taagcactgaggaagacagtgaggcggggcctccccgcga	2640		
tggggatggggtgcctggtagggcccccgccagtg	2680		
cacaccaggagattgaggaaaacttgcttagcttagagg	2720		
tggccatcagacaactggagggttggaagaggaattttg	2760		
tcgcctaagacccctcctgtcccagcttggggggactctg	2800		

FIGURE 15 cont.

24/34

2810 2820 2830 2840
tcccccaacctggctgcacctgaacgctctgctcagacag 2840
gcctttcatgaagagaagagtgggggagaggatgcagggg 2880
ggccccacccccacccccacagctgccacagcatctcacac 2920
cccagaggcctggagcagagggagattggctgaacttgat 2960
agcagacctgctcgggtccctgcctcccactctggcctct 3000
3010 3020 3030 3040
gcctttctccctcctgccttctgcttgggggaactcagggc 3040
ttcactcccagagcaccccccttaatccccacttttcaggc 3080
cgtctcagtattgcctatggggggtcacctccttccct 3120
accccaagtgcctttgctatgtttttatctcctggactgg 3160
aggtttatttttaatatatatattcttaagggaataaaaaaa 3200

FIGURE 15 cont.

SUBSTITUTE SHEET (RULE 26)

25/34

10	20	30	40
mddpsikkrilldlytdeheydevqei peessi appetsts 40			
htstngsshsgpgtatpggatsagpsagapqspv ivvds 80			
pel papkqksvknsskqkqqlankskiprpsl assls 120			
sl asslsghrdrkdrenqnavppqtppl ppsykqn 160			
qmngdstaaagggvsapatpttannnnashnngsimgggv 200			
210	220	230	240
qlnqsdnsnpvlqapgersslnitp lsrdisgghtqestt 240			
pattpstpslalpknfayltltvrkdsngygmkvsgdnpv 280			
fvesvkpggaeiaglvagdmilrvnghevrl ekhptvg 320			
likasttvelavkrsqkltrpssvsvtptstpi lsgdrdt 360			
asitgppqvdsikrremetykiqt lqkml eqeklnlerlk 400			
410	420	430	440
sdqnnpsyklseanirk lreqlhqvgaedaptvkl qaaag 440			
nkntalltpnqihlsasathsnqqfhh lhhhhnlhnnny 480			
ppqqqpastspafllsllprslssls lgrknktekdltts 520			
spfglttdflqqarmshqaesmsqsmhqhtstptsqqffh 560			
phqqqhrfketgptskgknkfllsrs l eedvppplpqrn 600			
610	620	630	640
pprqlnldlknagnaspaggshlvapvsdl dratspqlnrsg 640			
qqqlprstdnspnakskskiktkalsdpkmstqml lqm 680			
esasaagaaggsiev dggppplppr lpgmmedmsrgscq 720			
nlaqpnsvgtafnyp lvstttavqndnl niafplsqrpni 760			
vqqlaayqqaaqhamsggqatgal gqtpnl gknkhrrvgs 800			
810	820	830	840
spdnmhprhpdnritkttsgsweivekd gessppgtpplpy 840			
lssshmtvledpnennrgaaaagpgvfieshaftpmagas 880			
spipisihssmhhaaqsndtqkei ismedensdldepfid 920			
engpfnnltrlleaenvtflaif lnyvisnsdpapllfyl 960			
itelykegtskdmrkwaiyhstflvpraplswyrqdesl 1000			
1010	1020	1030	1040
arevdnvlqleydkveilrtvflrsrkrakdliseqiref 1040			
qakrtaglti ygptddkl aeaktdklreqi idkylmpnl 1080			
haliedengspdpvrkvalcsalstviyrifntrppps 1120			
ivervhhfvsrdksfksrimgknrk mnvrghplvlrqyye 1160			
vthcnhcqtiwgvspqgyhctdcklnihrqcskvvdsc 1200			
1210	1220	1230	1240
pgplpqaakrlahndkiskfmgk irprt sdvigne kr srq 1240			
eeldveltpdrgqasivrpsdrrpdanisisngntscn 1280			
tsglnttdlqssfhgscandsinpgggagcnmdlstsvas 1320			
ttpstsgsvaaglsafaelnaldtvdkearrerysqhpeh 1360			
ksapvsvnrse sykerlsknrrnsrrktsdpslssrpnd 1400			

FIGURE 16

26/34

1410	1420	1430	1440

eql d l g l s n a t y v g s s n s s l s s a g g s e s p s t s m e h f a a p g	1440		
a a g g v q v p p m g l n a n q h p h l l i q q h a q q y c q q d s f a a g l a	1480		
g a a g s s a a s n s s f w n a g h p l p v a r w t l e s e d e d d v n e a d w	1520		
s s m v a a e v l a a l t d a e k k r q e i i n e i y q t e r n h v r t l k l l	1560		
d r l f f l p l y e s g l l s q d h l l l l f p p a l l s l r e i h g a f e q s	1600		
1610	1620	1630	1640

l k a r r i e h n h v n t i g d l l a d m f d g a s g v l c e f a a q f c a	1640		
r a q i a l e a l k e k r n k d e m l a k l l k k s e s h k a c r r l e l k d l	1680		
l p t v l q r l t k y p l l f e n l y k v t v r l l p e n t t e a e a i q r a v	1720		
e s s k r i l l v e n q a v k t a e d a h k l a n i q r k l d r s s y d k e e f	1760		
k k l d l t q h h l l h d g n l t i k k n p s v a l h g l l f e n m i v l l t k	1800		
1810	1820	1830	1840

q d d k y y l k n l h t p l s i t n k p v s p i m s i d a d t l i r q e a a d k	1840		
n s f f l i k m k t s q m l e l r a p s s e c k t w f k h f s d v a a r a s k	1880		
n r s k n a s s n h d t s i s d p a l a a i p h s n t k e s l e l s t d t v a p	1920		
l a a t a t l t t t p l a p m l p i a t v t p a p a t n n s n v s s l t g v a l	1960		
r n p q r d a t a s e s d a d y v n t p k p r s s q n e v n r t m s i r s t g e	2000		
2010	2020	2030	2040

p i a k y s a n g t e a n d v t l r h s q s t r e s v r p g s t g e e r n s t y	2040		
g m v g g n s k r d s a s i v c s n n s n n t r t l l m a s p l v d p t a i q v	2080		
s i s p a h t a e p v l t p g e k l r r l d a s i r n d l l e k q k i i c d i f	2120		
r l p v e h y d q i v d i a m p e a p k d s a d i a l a a y d q i a t l t k m	2160		
l n e y m h v t p e q e v s a v s t a v c g h c h e k e k l r k k v a p s s s f	2200		
2210	2220	2230	2240

s s s p p p l p p p n r q h a q a q a q i p p s r l m p k l a t l d l d e v a i	2240		
h e d d d g y c e i d e l r l p a i p s k p h e r p t t p l a p f n t e p k t s	2280		
q s v i d a s k r a s t d a v p e g l l e a e p l e g d k t e t k g e d n e v k	2320		
t v p s d k l s e s c n e e r q c v e a d i t k e v a d p t t s k n e a a s v	2360		
d e l p s q s r e i k t a e n a s k s v a d k k e d n e e t i e e g v a s t v d	2400		
2410	2420	2430	2440

s s t a t s p t e s p k e t d k l t g g s s s t c g p n r i q h a s v l e p s v	2440		
p c h a l s s i v t i l n e q i s m l p k i n e r d m e r e r l r k e n q h l	2480		
r e l l s a l h d r q r v d e v k e t p f d l k k l m h a e d v e f d d d i d a	2520		
i s n s s l t p t p t p i p t a s p a s g v e t a e a m r i t s t e d e e	2559		

FIGURE 16 cont.

27/34

```

      10      20      30      40
      |      |      |      |
aagaacgtcgtgtgcgtgttgatggcgcgataagtgaaaa 40
aaataaaaacatTTTTcgtgtctttatggaatacaatacg 80
agtgcgtttttaagtTTAAGTgaaaagtagcagtgcaaaat 120
atatatacacatatcaaaacccattggaatagtgcAACCA 160
aaaaaacataaaacagaactaaagcaacgaaaatcgaggc 200
      210      220      230      240
      |      |      |      |
aattaaatcacttctggattcatttcggcccgTTTTtact 240
agctaattttaaaaatagctaccgaaagcattccgcaacca 280
cttctggtgactggccccacaccaatgcgaacaatactcc 320
acgggatcctccgacttcgagtggttacgcattgcctttg 360
aatttgcattcttcgcacccgcacagcagccagcaaaagcc 400
      410      420      430      440
      |      |      |      |
atcgcaggaaaccctatggatgaccatcaatcaaaaaaac 440
ggttactagatttatatactgacgaacatgaatacgatga 480
ggttcaagagataccagaagagtcgaagcattcagcccccg 520
gagacatcaacgagccataccagcacaaacggatccagcc 560
actcaggccctggaactgcaactggaccaggagcgcacatc 600
      610      620      630      640
      |      |      |      |
tgcaggcccgctcagcaggtgcgccacagtcaccagtaatt 640
gttgtggactcgggtcccgagctgccagcgcaccaagcaga 680
aatctgttaagaactcgaagagcaaacagaaagcaaaagca 720
gttggcgaacaaatcgaagattccacgatctccttcgttg 760
gcgagcagcctgagtagtctggccagcagccttagtggtc 800
      810      820      830      840
      |      |      |      |
atagggatcgggacaaagatcgggacaaggatcgggagaa 840
ccagaacgccgtgccgcgcagacgccgccgttgccaccg 880
agctacaagcagaaccaaatgaatggtgactctacggctg 920
cagctggtggtggtgtttctgccccagccacgcccaccac 960
cgccaataataacaatgcgagccacaacaacggcagcata 1000
      1010      1020      1030      1040
      |      |      |      |
atgggcggaggcgtgcaatigaatcaatcggacaactcca 1040
accccggttctccaggcgccgggggagcgcagtagcctcaa 1080
tctgactccctctcccggtatctgagtggtggccacacc 1120
caggagtcacagacgccagcgaaccacgccgagtagcggaa 1160
gcctagcattaccaaaaaattttcagtatttaaccctgac 1200
      1210      1220      1230      1240
      |      |      |      |
tgtgcgaaaagatagcaacgggatacgggatgaagggtttcc 1240
ggagataatcctgtgtttgtggagagcggttaaacccggag 1280
gcgcagcggagattgcaggcctggttgctggcgatatgat 1320
actaagggttaaacggccatgaagtgcgactagagaagcat 1360
ccaactgtagtgggtcttataaaagcctcgacaactgttg 1400

```

FIGURE 17

28/34

1410	1420	1430	1440
agctggcgggtgaagcgaagtcagaagctaacgcgaccctc	1440		
ttcgggtatcagtggtaacgccctcgacacccattctctct	1480		
ggacgagatcgtaccgcttcaataaactgggccacagccgg	1520		
tggacagtattaaacgaaggagatggagacttacaaaat	1560		
tcagacggttgcaaaaaatgctggagcaggagaaactaaat	1600		
1610	1620	1630	1640
ctggagcgaattaaaaagcgnacaaaataatccgagctaca	1640		
agctatctgaggcgaatatccgtaagctgcgcgagcaact	1680		
tcatcaagtgggagctgaggatgcaccgactgttaaactt	1720		
caggcggccgctggcaataagaacacagcattactaacgc	1760		
ccaaccaaatccaacacttgctccgcgtccgccactcacag	1800		
1810	1820	1830	1840
caatcagcaattccatcatcttcaccaccaccacaatctc	1840		
cacaacaacaattatccgccacagcaacagccagctagca	1880		
cctcaccgcattcctgtccctcctgccgcgttccctttc	1920		
gtccttgctcgctgggcacgcgcacaaaacaagaccgaaaag	1960		
gacctaacgacttcatctccctttggcctaacgacggatt	2000		
2010	2020	2030	2040
tcctgcagcagcaacggatgagccaccaagcggagtcgat	2040		
gtctcaatcgatgcatacagcacaccagcactccgacctcg	2080		
cagcagttcttccatcgcatacagcaacaacatcgcttta	2120		
aggaaactggaccgacgtcgaaaggcaagaacagttcct	2160		
aatttcgaggagtttgatcgaggaggatgtgccgccacca	2200		
2210	2220	2230	2240
ctgccgcagaggaatccgccccaggcagttaaatttggacc	2240		
tgaagaacggaaacgcgtcgccgggtgggtcacatctagt	2280		
ggctccagtttccgatctggatcgccgccacaagtccccaa	2320		
ttaaatagatcccaacaacagcaactgcctaggagcactg	2360		
acaacagccccgagcaatgccaaagtccaagcgttccaagat	2400		
2410	2420	2430	2440
caaaacgaaggccctatcggtatcctaagatgtccactcag	2440		
atgctcctgcbaatggaatcggcgagtgacgctggagcag	2480		
ccggagggttccatcgaggtagatgggggtccaccaccgct	2520		
accgcctcgcttgccctggcatgatgacggaggatgatgagc	2560		
cgcggcagctgtcagaatctcgctcaacccaattccgtgg	2600		
2610	2620	2630	2640
gcactgccttcaactatcctctggtgtccacaacaacagc	2640		
ggtgcagaacgataacctgaacattgcctttcctttgtcc	2680		
caacgaccaaacattgtccagcagctgcagcaatatcaac	2720		
agcagcagcagcatcaaatgagcgggtggccagggccacagg	2760		
cgctctgggacagactccgaatttgggaaaaaacaagcat	2800		

FIGURE 17 cont.

29/34

2810	2820	2830	2840
cgcgcggttggttcttcaccggacaatatgcacccacgtc 2840			
atccagatcggataacgaaaacgacttcgggctcgtggga 2880			
gattgttgaaaaggatggcgaatcctccccgcccggaaca 2920			
ccgccgttccatattctatccagctcccacatgaccgtgc 2960			
tggaagatccgaatgagaacaatcgtggagcagcagcagc 3000			
3010	3020	3030	3040
cggacctggagtcctttatcngtgcacacagtttacgccg 3040			
atggcgggagcctcttctccgatcccgatattccttacatt 3080			
ccagccacatgcacgcggcccagtcgaacgatagcagaa 3120			
ggagatcatctcgatggaggacgaaaactcggacttgga 3160			
gagcccttcattgacgagaacggaccctttaacaattctaa 3200			
3210	3220	3230	3240
ctcgtttgttagaggccgagaacgtcactttcctagccat 3240			
cttccctaaactacgtgatctcaaaactcggatcccgcgcca 3280			
cttctgttttacctgattactgagttgtacaaggaggaggca 3320			
cctccaaggacatgcggaaatgggcctacgaaatccactc 3360			
cacattcctcgtgccgcgggctccattgtcatggtatcgc 3400			
3410	3420	3430	3440
caagatgaatcgctggcccgcgaggtggataatgtccigc 3440			
agttggagtatgacaaagtggagatcctaaggacagtttt 3480			
tctgcgtagccgaaagcgggccaaggacctaatacgtgag 3520			
cagctgcgtgagtttcagcagaagcgcaccgcggcctgg 3560			
gaaccatttacggacccacggacgacaagctggccgaggc 3600			
3610	3620	3630	3640
gaagacggataagctaaggaggacaaatcatcgacaaatat 3640			
ttaatgcccaatcttcacgcgcttattgaggatgagaacg 3680			
gttcaccgcccggaggatgtacgcaagggtggcgttgtgttc 3720			
tgctctttccaccgtcatctaccgtattttcaacactcgt 3760			
ccgcctccaagcagcatcggttgagcgggtccatcactttg 3800			
3810	3820	3830	3840
tgagcagggacaagagtttcaagtcgcgtattatgggcaa 3840			
aaatcgcaagatgaatgttcgtggatcatccattggtattg 3880			
cgtcaatactatgaagtgcgcactgcaatcattgtcaga 3920			
cgattatctggggcgtgagcccgcgaaggttatcattgtac 3960			
agactgtaaatggaacatacacccgtcagtgctcgaagta 4000			
4010	4020	4030	4040
gtggacgagagttgtcccgggtcccttgccccaggcaaaac 4040			
gtctcgcccacaacgacaagatcagtaaatcatgggcaa 4080			
aattcgaccgcgtaccagcgacgtcattggaaatgaaaag 4120			
cgaagtcgtcaagatgaggaattggatgttgagttgactc 4160			
cagaccgtggtcaggcgtcgattgtgcgtcaaccctctga 4200			

FIGURE 17 cont.

SUBSTITUTE SHEET (RULE 26)

4210 4220 4230 4240

4410 4420 4430 4440

4610 4620 4630 4640

4810 4820 4830 4840

5010 5020 5030 5040

5210 5220 5230 5240

5410 5420 5430 5440

SUBSTITUTE SHEET (RULE 26)

31/34

5610	5620	5630	5640
.....			
aggtcaaccaggcagtaaaagacagcagaggatgctcaca			5640
gctgcaaaacattcagcgttaagttagacagatcctcctat			5680
gacaagggaagagtttaagaaattggacctgacccaacatc			5720
acctcatccatgacggcaatctgacgatcaagaagaatcc			5760
tagcgtgcagctacatggacttctgtttgaaaacatgatt			5800
5810	5820	5830	5840
.....			
gttttgctgaccaagcaggatgataaatattatctaaaga			5840
acttgcacaccccgcctatcgatcaccaataagccagtcag			5880
tccaattatgagcattgatgaggacactttgatccggcag			5920
gaggcggctgataaaaatttccttttctcatcaagatga			5960
agacatcacaaatgttggagctacgcgcgcctagtagctc			6000
6010	6020	6030	6040
.....			
ggagtgcaagacatggtttaaacacttctcggtatgtggct			6040
gctcgtcagtcctaagaatcggttcaaagaacgcataagca			6080
accatgacacgagttatagtgatccagctctcgccgctat			6120
tcgcgattccaacaccaaagagtcgttggagttgagcact			6160
gatcacgtacagccattggctgcgacagccacattgacta			6200
6210	6220	6230	6240
.....			
ccacaccattggccccaatgctgcctatagccacggttac			6240
accggctccagcgaccaataatagtaacgttagctctctt			6280
actggagttcagttgcgaaccctcaacgggatgagcag			6320
caagtgaatctgatgaggattatgtaaacacaccaaagcc			6360
gcgttcgagccaaaatgaagttaatcgcaactatgtccata			6400
6410	6420	6430	6440
.....			
agaagcactggcgaacccattcagaagtattcggcgaatg			6440
ggacggaagcaaacgacgttactttacgacactctcagtc			6480
gactaggggaatcggttagaccaggatctactggggaggag			6520
cgaaactccacgtatggtatggttggaggtaactccaaac			6560
gcgacagcgccagcattgtctgctcgaacaactcgaacaa			6600
6610	6620	6630	6640
.....			
cacgcgcacccttctgatgcagagtccttgggtggaccct			6640
acggccattcaggtcagcattagtcgctcacacagcgg			6680
aacctgtgttgacaccaggagagaagttaacgccgcttggg			6720
cgcctccattaggaatgatttgcctggagaagcagaaaatc			6760
atttgtgatatcttccgtttgccagtggaaacactacgacc			6800
6810	6820	6830	6840
.....			
agattgtggacattgccatgatgccagaggcgccgaaaga			6840
cagtgacagatattgcttttagctgcttacgatcagattcaa			6880
accctgaccaagatgctgaacgagtacatgcacgtcacgc			6920
ctgagcaagaggctcagcggtgtccacggcggtttgtgg			6960
ccactgtcacgagaaggaaaagcttcgtaagaagggtggca			7000

FIGURE 17 cont.

32/34

7010	7020	7030	7040
ccatcctcatcggttttcttcatcaccaccaccgctgccgc 7040			
ctcccaataggcagcatgccagggctcaggcccagatacc 7080			
gccatcgcggtgatgcccaactacaaactcttgatctt 7120			
gacgaagttagccatacacgaagacgatgacggatactgtg 7160			
agatcgacgaactgcgcttacccggctattccgtccaaacc 7200			
7210	7220	7230	7240
acatgagcggcccacaaacgccactggctcctttcaatact 7240			
gagccgaaaacttcacaatctgttatagatgcctcgaaac 7280			
gtcaatccactgatgccgttcggagggttactggaaca 7320			
agaaccactcgaaggcgataagacggagaccaagggtgaa 7360			
gataatgaagtgaaaactgtgccgtcagataagctaagtg 7400			
7410	7420	7430	7440
aatcatgcaatgaagagaggcaatgtgtggaggcggtat 7440			
cacaaggaagtggcagatccaacgacctctaagaatgaa 7480			
gctgcagcatcggtggatgaattaccaagccagagccggg 7520			
agataaaaacggctgaaaacgcaagcaaatctgtagctga 7560			
caaaaaggaagacaacgagggaaccatcgaagaagggtgtg 7600			
7610	7620	7630	7640
gcattcacggtcgatagctccactcaaacatcaccaactg 7640			
aatcgccgaaagagccggataagtttaactggaggatcgag 7680			
cagcacctgtgggcccgaatcgcatcagcacgctagtgtg 7720			
ctggagccgagtgtgccctgccatgcactcagcagcattg 7760			
taacaatactgaatgagcagatttccatgcttttgccaaa 7800			
7810	7820	7830	7840
aattaacgaacgcgatatggaaaggagcgattgcgtaaa 7840			
gagaatcaacaccttcgcgagctcttgagtgcgctgcctg 7880			
atcgacagcgagttgatgaagttaaaggaaactccgtttga 7920			
tctaaagaagctgatgcattgctgaggatgtagagtttgac 7960			
gatgatattgacgccatttccaacagttcgtgacgccaa 8000			
8010	8020	8030	8040
cgctacgccgattcccacagcatcaccaagcgccagcgg 8040			
ccaggtagagacagcgggaagccatgaggattactagcaact 8080			
gaggatgagggaataggaaacttgtttaatttttttttgta 8120			
taaaatacatattttgttcttggtttttcccgcacgcatttg 8160			
accaattgaattggcggcgtctatgtttttacgagtagcct 8200			
8210	8220	8230	8240
ttcttagagcatgccttttactactgttaatgtagcttca 8240			
tttgttttcttaaaatattatgtattccacaagagccgat 8280			
aacacttttaaatgaagtctaacattgtccccctcgtaaaac 8320			
taattaaatacatatatatatgcatttatcgacttaaaa 8360			
aatatcagataaacacacaatgtttgatataattataaag 8400			

FIGURE 17 cont.

33/34

8410 8420 8430 8440
aacaataaacaacatagttattgagaagaatagcaaaaaa 8440
aaaataaaaaaaaaaaaaaaaaaaaaa 8464

FIGURE 17 cont.

SUBSTITUTE SHEET (RULE 26)

SUBSTITUTE SHEET (RULE 26)

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/12, C07K 14/82, A61K 38/17, G01N 33/68, C12N 15/63, 5/10	A3	(11) International Publication Number: WO 99/47557 (43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/US99/06051 (22) International Filing Date: 18 March 1999 (18.03.99) (30) Priority Data: 60/078,634 18 March 1998 (18.03.98) US (71) Applicant: ONYX PHARMACEUTICALS, INC. [US/US]; 3031 Research Drive, Richmond, CA 94806 (US). (72) Inventors: BOLLAG, Gideon; 172 Catalina Drive, Hercules, CA 94547 (US). HART, Matthew, J.; 1210 Masonic Avenue, Berkeley, CA 94706 (US). ROSCOE, William; 3099 California Street, San Francisco, CA 94115 (US). POLAKIS, Paul; 509 Barone Lane, Mill Valley, CA 94941 (US). STERNWEIS, Paul; 2103 Flat Creek Drive, Richardson, TX 75080 (US). KOZASA, Tohru; 7415 Centenary, Dallas, TX 75225 (US). JIANG, Xuejun; 2400 Waterview Parkway #932, Richardson, TX 75080 (US). (74) Agent: GIOTTA, Gregory; Onyx Pharmaceuticals, Inc., 3031 Research Drive, Richmond, CA 94806 (US).	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 18 November 1999 (18.11.99)	
(54) Title: IDENTIFICATION OF FACTORS WHICH MEDIATE THE INTERACTION OF HETEROTRIMERIC G PROTEINS AND MONOMERIC G PROTEINS (57) Abstract Monomeric GTPase guanine nucleotide exchange factor (GEF) have been identified which also contain an RGS region analogous to those of GTPase activating proteins (GAP). One of these GEF proteins, a Rho GEF has been demonstrated to contain an RGS sequence that has GAP activity toward a α subunit of a heterotrimeric G prote in.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/06051

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/82 A61K38/17 G01N33/68 C12N15/63
C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MATTHEW J. HART ET AL.: "Identification of a novel Guanine nucleotide exchange factor for the Rho GTPase" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 41, 11 October 1996 (1996-10-11), pages 25452-25458, XP002057776 MD US cited in the application the whole document --- -/--	1-28

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

21 September 1999

Date of mailing of the international search report

05/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Montero Lopez, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/06051

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DAVID M. BERMAN ET AL.: "GAIP and RGS4 are GTPase-activating proteins for the Gi subfamily of G protein alpha subunits" CELL, vol. 86, no. 3, 9 August 1996 (1996-08-09), pages 445-452, XP002111126 NA US cited in the application the whole document ---	1-28
P,X	KOZASA T ET AL: "p115 RhoGEF, a GTPase activating protein for Galpha12 and Galpha13 'see comments!'" SCIENCE, (1998 JUN 26) 280 (5372) 2109-11. , XP002111127 abstract page 2109, right-hand column, paragraph 2 -page 2111, middle column, paragraph 2 ---	1-28
P,X	ALAN HALL: "G proteins and small GTPases: Distant relatives keep in touch" SCIENCE, vol. 280, 26 June 1998 (1998-06-26), pages 2074-2075, XP002111128 DC page 2074, middle column, paragraph 2 -page 2075, right-hand column, paragraph 2 ---	1-28
P,X	HART M J ET AL: "Direct stimulation of the guanine nucleotide exchange activity of p115 RhoGEF by Galpha13 'see comments!'" SCIENCE, (1998 JUN 26) 280 (5372) 2112-4. , XP002111129 the whole document ---	1-28
P,X	MAO J ET AL: "Guanine nucleotide exchange factor GEF115 specifically mediates activation of Rho and serum response factor by the G protein alpha subunit Galpha13." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1998 OCT 27) 95 (22) 12973-6. , XP002111130 abstract page 12974, right-hand column, paragraph 2 -page 12976, right-hand column, paragraph 1 --- -/--	1-29

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/06051

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 98 20127 A (ONYX PHARMACEUTICALS) 14 May 1998 (1998-05-14) page 2, line 13 -page 9, line 32 page 19, line 13 -page 23, line 2 page 36, line 8 -page 37, line 32 ----	19-25
P, X	WO 98 57990 A (ONYX PHARMACEUTICALS) 23 December 1998 (1998-12-23) page 3, line 16 -page 4, line 2 page 4, line 19 -page 5, line 13 page 6, line 20 -page 7, line 3 page 8, line 26 -page 9, line 12 page 19, line 14 -page 20, line 11 page 20, line 24 -page 24, line 30 ----	19-25
T	FUKUHARA S ET AL: "A novel PDZ domain containing guanine nucleotide exchange factor links heterotrimeric G proteins to Rho." JOURNAL OF BIOLOGICAL CHEMISTRY, (1999 FEB 26) 274 (9) 5868-79. , XP002111131 abstract page 5873, left-hand column, last paragraph -page 5874, right-hand column, paragraph 3 page 5875, right-hand column, paragraph 2 -page 5878, right-hand column, last paragraph -----	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/06051

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 18
is directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/06051

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9820127	A	14-05-1998	AU	4801097 A	29-05-1998
WO 9857990	A	23-12-1998	AU	7966498 A	04-01-1999

